



# Molecular Tools for the Detection and Deduction of Azole Antifungal Drug Resistance Phenotypes in *Aspergillus* Species

Anna Dudakova,<sup>a</sup> Birgit Spiess,<sup>b</sup> Marut Tangwattanachuleeporn,<sup>a,c</sup> Christoph Sasse,<sup>d</sup>  
 Dieter Buchheidt,<sup>b</sup> Michael Weig,<sup>a</sup> Uwe Groß,<sup>a</sup>  Oliver Bader<sup>a</sup>

Institute for Medical Microbiology, University Medical Center Göttingen, Göttingen, Germany<sup>a</sup>; 3rd Department of Internal Medicine, Hematology and Oncology, Mannheim University Hospital, University of Heidelberg, Mannheim, Germany<sup>b</sup>; Unit of Medical Technology, Faculty of Allied Health Sciences, Burapha University, Chon Buri, Thailand<sup>c</sup>; Institute of Microbiology and Genetics, Department of Molecular Microbiology & Genetics, Georg August Universität Göttingen, Göttingen, Germany<sup>d</sup>

<b>SUMMARY</b> .....	<b>1065</b>
<b>INTRODUCTION</b> .....	<b>1066</b>
<b>SCREENING FOR AZOLE-RESISTANT <i>ASPERGILLUS</i> STRAINS</b> .....	<b>1069</b>
<b>SPECIES IDENTIFICATION OF <i>ASPERGILLUS</i></b> .....	<b>1073</b>
<b>SUSCEPTIBILITY TESTING PROCEDURES FOR <i>ASPERGILLUS</i> SPP.</b> .....	<b>1073</b>
<b>EVOLUTION OF <i>cyp51</i> GENE FAMILIES</b> .....	<b>1074</b>
<b>RESISTANCE THROUGH TRANSCRIPTIONAL MODULATION OF <i>cyp51</i> GENES</b> .....	<b>1075</b>
<b>RESISTANCE-CONFERRING ALTERATIONS WITHIN <i>Cyp51</i> PROTEINS</b> .....	<b>1076</b>
<i>Aspergillus fumigatus</i> .....	<b>1076</b>
<i>Aspergillus flavus</i> .....	<b>1078</b>
Other <i>Aspergillus</i> Species .....	<b>1080</b>
<b>MOLECULAR METHODS FOR DETECTION OF ALTERATIONS AT <i>cyp51</i> LOCI</b> .....	<b>1080</b>
<i>Aspergillus fumigatus</i> .....	<b>1080</b>
Other <i>Aspergillus</i> Species .....	<b>1082</b>
<b>OTHER RESISTANCE MECHANISMS</b> .....	<b>1082</b>
<b>CONTRIBUTION OF PHYLOGENETIC ANALYSES TO RESISTANCE DETECTION</b> .....	<b>1083</b>
<b>CONCLUSIONS</b> .....	<b>1083</b>
<b>ACKNOWLEDGMENTS</b> .....	<b>1084</b>
<b>REFERENCES</b> .....	<b>1084</b>
<b>AUTHOR BIOS</b> .....	<b>1090</b>

Published 13 September 2017

**Citation** Dudakova A, Spiess B, Tangwattanachuleeporn M, Sasse C, Buchheidt D, Weig M, Groß U, Bader O. 2017. Molecular tools for the detection and deduction of azole antifungal drug resistance phenotypes in *Aspergillus* species. Clin Microbiol Rev 30:1065–1091. <https://doi.org/10.1128/CMR.00095-16>.

**Copyright** © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Oliver Bader, [oliver.bader@med.uni-goettingen.de](mailto:oliver.bader@med.uni-goettingen.de).

**SUMMARY** The incidence of azole resistance in *Aspergillus* species has increased over the past years, most importantly for *Aspergillus fumigatus*. This is partially attributable to the global spread of only a few resistance alleles through the environment. Secondary resistance is a significant clinical concern, as invasive aspergillosis with drug-susceptible strains is already difficult to treat, and exclusion of azole-based antifungals from prophylaxis or first-line treatment of invasive aspergillosis in high-risk patients would dramatically limit drug choices, thus increasing mortality rates for immunocompromised patients. Management options for invasive aspergillosis caused by azole-resistant *A. fumigatus* strains were recently reevaluated by an international expert panel, which concluded that drug resistance testing of cultured isolates is highly indicated when antifungal therapy is intended. In geographical regions with a high environmental prevalence of azole-resistant strains, initial therapy should be guided by such analyses. More environmental and clinical screening studies are therefore needed to generate the local epidemiologic data if such measures are to be implemented on a sound basis. Here we propose a first workflow for evaluating isolates from screening studies, and we compile the MIC values correlating with individual amino acid substitutions in the products of *cyp51* genes for interpretation of DNA sequencing data, especially in the absence of cultured isolates.

**KEYWORDS** *Aspergillus fumigatus*, *Aspergillus flavus*, *Aspergillus terreus*, *Aspergillus niger*, azole drug resistance, diagnostics, *cyp51A*, *cyp51B*, *cyp51C*, efflux, tandem repeats

## INTRODUCTION


Inhalation of *Aspergillus* spores, primarily of the species *Aspergillus fumigatus*, might result in different clinical manifestations. Whereas sequelae are usually absent in a healthy host, frequent exposure to spores might foster the development of allergic bronchopulmonary aspergillosis in individuals with an atopic disposition. Cavitary lung disease, e.g., following tuberculosis, might predispose individuals to aspergilloma. Chronic necrotizing aspergillosis is predominantly observed in patients who suffer from chronic lung disease or mild immunosuppression. Invasive pulmonary aspergillosis is the most dangerous and life-threatening clinical manifestation and might result from *Aspergillus* exposure in heavily immunosuppressed patients (1). In fact, *Aspergillus* spp., primarily *A. fumigatus*, are the most common emerging fungal pathogens, especially in patients with malignant hematological diseases who are undergoing intensive chemotherapy or after allogeneic hematologic stem cell transplantation. Consequently, prolonged neutropenia ( $<500$  cells/ $\mu$ l for more than 10 days) has been identified as a major risk factor (1).

Over time, several azole-based antimycotic drugs have been developed, among which itraconazole (ITZ), voriconazole (VRZ), posaconazole (PSZ), and (lately) isavuconazole (ISAZ) are mainly used for the treatment and/or prophylaxis of aspergilloses (Fig. 1). Azoles are steric inhibitors of sterol 14 $\alpha$ -demethylase enzymes catalyzing a critical step in ergosterol biosynthesis (Fig. 2A and B).

The infection-related mortality rates are still exceptionally high, despite recent improvements in prophylaxis, early diagnosis, and antifungal treatment of the disease (2). The incidence of azole resistance in *Aspergillus* spp. has increased over the past years, additionally jeopardizing the outcomes for high-risk populations by failure of azole-based prophylaxis and first-line treatment of invasive aspergillosis.

Starting with the initial observation in a Dutch strain collection that over the 1990s numbers of azole-resistant *A. fumigatus* (ARAF) had increased unproportionally due to a single resistance allele (3), the same has now been demonstrated on a global level (4). Different patterns of resistance are seen, with multiazole and pan-azole resistance being more common than resistance to a single triazole. A variety of mutations in the coding region of *cyp51A* (Fig. 2C), a gene encoding a sterol 14 $\alpha$ -demethylase in *A. fumigatus*, as well as tandem repeats of 34, 46, and 53 bp upstream in its promoter region (Fig. 2E), have been found to confer, or at least to correlate with, various degrees of drug resistance. Today, the most frequently observed resistance allele consists of a 34-bp tandem repeat in the promoter region of *cyp51A* combined with the L98H substitution (TR<sub>34</sub>/L98H). This allele confers high ITZ resistance and various levels of cross-resistance to other azoles. Together with strains carrying a TR<sub>46</sub> allele in conjunction with Y121F and T289A exchanges, such strains are thought to spread globally through the environment (5), potentially propagated by the use of 14 $\alpha$ -demethylase inhibitor (DMI) fungicides in agriculture. In fact, for many plant-pathogenic fungi, similar resistance mechanisms toward agricultural azole-based fungicides are observed (6–10), and their spread to the level of complete replacement of susceptible strains in some regions (11, 12) is a much discussed topic in phytopathology. The best-studied example is the wheat pathogen *Zymoseptoria tritici*, in which the *cyp51B* paralog has evolved several resistance mutations in both promoter and coding regions as a result of azole selection (for an overview, see reference 12).

Resistant *Aspergillus* strains present in the environment are thought to exogenously colonize or infect susceptible hosts. These observations are of significant concern, as exclusion of azole-based antifungals from prophylaxis or first-line treatment of invasive aspergillosis dramatically limits treatment options (4). A careful reevaluation of remaining management options for invasive aspergillosis caused by azole-resistant strains by








**FIG 1** Chemical structures of clinically used azole antifungals. Azoles are characterized by five-atom heterocycles which contain at least one nitrogen atom (red). Compounds containing moieties with two nitrogen atoms are called diazoles, and those with three nitrogen atoms are called triazoles. Recently marketed antifungals contain one (B and D) or more (A, C, E, and F) triazole moieties and a benzene ring substituted with fluorine (A to C and F) rather than chlorine (D and E). Triazole antifungals are derivatives of either fluconazole (A to C) or ketoconazole (D to F) as the lead compound. This correlates with cross-resistance phenotypes observed in clinical and environmental isolates (see the text).

an international expert panel suggested that, in initial therapy, the environmental prevalence of azole-resistant strains should be considered in areas where the prevalence reaches 10% (13, 14). For example, this may be done by combining VRZ with an echinocandin or liposomal amphotericin B (13, 15).

In about 20 to 50% of azole-resistant clinical *A. fumigatus* isolates, no mutations within the *cyp51A* locus are observed (16–18); therefore, additional pathways and factors that confer secondary resistance must exist. In this context, only the HAP complex member *hapE* (CCAAT binding complex [CBC]) has been implicated in clinical ARAf so far (19, 20). Additionally, drug efflux has been discussed to contribute to resistance (17, 21, 22). Hence, different azole resistance patterns may be multifactorial, and this must also be considered when encountering isolates with an altered *cyp51* sequence (23).

The molecular detection of genetic alterations leading to azole resistance is gaining importance, given the poor culture-based diagnostic yield for clinical specimens from

**A** Normal susceptible cell in absence of azoles**B** Normal susceptible cell in presence of azoles**C** Mutations in *cyp51A***D** Overexpression of *cyp51A***E** Increased expression of efflux pumps

**FIG 2** Azole antifungal drug resistance mechanisms in fungal cells. (A) *A. fumigatus*, including the cell wall (CW) and the cytoplasmic membrane (CM), in the absence of azoles. Cyp51 activity is required for the biosynthesis of the membrane compound ergosterol (green). The *abcA*, *cdr1B*, and *mdr1* genes encode efflux pumps, which are localized in the plasma membrane. These genes are regulated by different transcription factors (TF), such as AtrR. (B) In the presence of azoles, some drug molecules can be pumped out through efflux pumps, but intracellular levels are sufficient to inhibit Cyp51A, resulting in decreased amounts of ergosterol in the membrane. (C) Mutations in *cyp51A* (asterisk) can reduce target binding of the antifungal drug, therefore conferring resistance. (D) Overexpression of the drug target Cyp51A can be mediated by mutations in the HAP complex (HAP\*) or by different kinds of tandem repeats within the promoter region of *cyp51A*. (E) Increased expression of efflux pumps, such as *AbcA*, *Cdr1B*, and *Mdr1*, increases the drug tolerance of *A. fumigatus*. Possible mechanisms for increased transcription may be gain-of-function mutations in the regulating transcription factors (TF\*, AtrR\*).

hematological patients. From a clinical point of view, azole resistance is associated with therapeutically relevant azole treatment failure that is increasingly reported and plays a more and more important role, especially in hematology and intensive care unit (ICU) patients, leading to high mortality rates (24–27).

There is an ongoing debate as to how epidemiologic thresholds should be generated and to what degree laboratory findings of azole resistance in *A. fumigatus* outside invasive aspergillosis should be reported to clinicians (28). Nevertheless, it is evident that medium- to large-scale environmental and clinical screening studies are urgently needed to generate epidemiologic data to reassess clinical treatment options on local or national levels. Molecular analysis of such isolates is not readily available to most diagnostic laboratories and is far from being standardized. Therefore, we outline here a stepwise workflow for evaluating isolates from such screening studies to build up local epidemiologic data (Fig. 3). Furthermore, we created an annotated reference sequence for easy identification of known polymorphisms in *A. fumigatus* Cyp51A, in accordance with their recently proposed unified nomenclature (29), and compiled the MIC values correlating with individual Cyp51A amino acid substitutions. This should aid in the interpretation of DNA sequencing data, especially in the absence of a culture isolate.

### SCREENING FOR AZOLE-RESISTANT *ASPERGILLUS* STRAINS


Clinical guidelines, e.g., by the Infectious Diseases Society of America (IDSA) (30, 31), do not specify laboratory testing procedures but rather reference the general literature on the isolation of *Aspergillus* from respiratory tract specimens (32, 33).

Only recently has it been recommended “. . .to test different, up to five, colonies as different azole susceptibility phenotypes might be present in a single culture” (13). Indeed, rapid conidial dispersion is a key feature of *Aspergillus* spp., especially *A. fumigatus* (34). This allows the fungi to rapidly colonize fresh materials and, presuming a certain selective pressure in the environment, also explains the rapid clonal expansion of resistance phenotypes (35) and makes mixed samples very likely. Dissemination of resistant isolates is also directly propagated by human activities (36).

A very educative experiment highlighting how easily conidia of *A. fumigatus* are moved through the air was conducted by Kwon-Chung and Sugui (34). In their setup, fresh agar plates were kept next to a sporulating *A. fumigatus* culture plate. Compared to growth in the same setup with *A. nidulans*, growth was observed on the fresh plates in a very rapid fashion, despite all attempts to keep the environment free of air movement. The lecture to be taken here with respect to laboratory work is that, when handling aspergilli, especially *A. fumigatus*, working in the best possible sterile environment free of air movement is highly advisable to avoid cross-contamination between samples and/or cultures. In addition, both environmental and clinical samples may contain susceptible along with resistant fungal material (37, 38). This leads to a reduced detection rate for resistant strains under regular culture conditions in cases where susceptible fungal cells pose the majority of the inoculum. In such cases, additional use of screening agars with included antifungal drugs may be highly beneficial to suppress the growth of most susceptible strains (Fig. 3). For some samples, this may lead to the detection of a single resistant colony only. This emphasizes the need to reduce the number of cross-contaminating conidia from previously cultured samples. Best practice will physically separate the primary culture from any subsequent handling of cultures, especially those with resistance phenotypes.

Standardized protocols for screening of environmental samples do not yet exist. This may be due partially to the heterogeneity of such “environmental samples,” which may encompass dry soil (e.g., see references 39 to 43), air samples (35, 44–46), plant material (43), or simply surface swabs (46). For soil, rigorous vortexing in NaCl (43) and plating of the supernatant following settling of the soil debris have been used. Addition of 0.5% (wt/vol) saponin facilitated the extraction of highly hydrophobic material, i.e., *A. fumigatus* conidia (39, 40). To restrict the growth of most potential contaminants, incubation at elevated temperatures (>43°C) can be used, as *A. fumigatus* is highly





**FIG 3** Potential sampling workflow for ARAf screening studies. No standardized scheme for conducting screening studies is established yet, but combining several approaches proposed in the literature gives rise to an efficient workflow that eliminates false-positive results and yields robust numbers on the prevalence and phylogenetic cohesion of resistant isolates.

thermotolerant (47). However, especially environmental samples from tropical areas may still yield strong overgrowth with other thermotolerant species (our own unpublished observation).

Drug concentrations for screening agar plates used in the literature range from 0.5  $\mu\text{g/ml}$  for PSZ (48, 49) to 1 to 4  $\mu\text{g/ml}$  for ITZ or VRZ (39, 43, 48–50). At the lower end of the concentration range, a higher false-positive rate may be observed (39). When working with the goal of isolating *A. fumigatus*, specifically screening for fluconazole resistance is not necessary, as *A. fumigatus* is intrinsically resistant to this compound (51–53).

Different mutations leading to singly elevated resistances to particular drugs only are known (Tables 1 and 2). MIC values measured for isolates with altered azole susceptibility are cumulated in Tables 1 and 2. MIC values for tests done according to

**TABLE 1** Levels of drug resistance conferred by mutations in *A. fumigatus cyp51A*, using EUCAST breakpoints<sup>a</sup>

Mutation	No. of isolates with MIC (μg/ml)															
	Itraconazole								Voriconazole							
	S	I	R	S	I	R	S	I	R	S	I	R	S	I	R	S
TR <sub>34</sub> /L98H				7	4	18	13	32	13	55	7	7				
TR <sub>34</sub> /L98R																
TR <sub>34</sub> /L98Y										1						
TR <sub>34</sub> /L98Q										1						
TR <sub>34</sub> /L98I										1						
TR <sub>34</sub> /L98H/S297T/F495I										6						
TR <sub>46</sub> /Y121F/T289A	5	2	2							1						
TR <sub>46</sub> /Y121F/T289A/G448S										1						
TR <sub>46</sub> /Y121F/T289A/G448S										1						
TR <sub>46</sub> /X3/Y121F/M172I/T289A/G448S										1						
F46V/M172V/N248T/D255E										1						
G54E										8						
G54R										1		1				
G54R/M220T										2		2				
G54W										1		1				
Y121F	1									2		1				
P216L										1		4				
F219C										1		1				
M220I										6		1				
M220K										3		1				
M220L										3		1				
M220T										3		1				
M220V										1		1				
M220R										1		1				
Y431C										2						
G432A										1		1				
G434C										1		1				
G448S										2		1				

<sup>a</sup>Values are cumulative numbers of isolates with the respective MIC values reported in the literature, stratified according to clinical EUCAST breakpoints (133). S, susceptible; I, intermediate; R, resistant.

TABLE 2 Levels of drug resistance conferred by mutations in *A. fumigatus cyp51A*, using CLSI breakpoints<sup>a</sup>

No. of isolates with MIC (μg/ml)																																						
Itraconazole																Posaconazole																						
Voriconazole																Posaconazole																						
S		I		R		S		I		R		S		I		R		S		I		R		S		I		R										
<1	1	2	4	>4	8	>8	16	>16	32	>32	<0.125	0.125	0.25	0.5	1	2	>2	4	>4	8	>8	16	>16	<0.125	0.125	0.25	0.5	1	2	4	>4	8	>8	16	>16	>32		
Mutation																																						
TR <sub>34</sub> /L98H	11				2	14	12	59							5	28	5	24	23	6	7	5			1	3	50	21	12	2								
TR <sub>46</sub> /Y121F/T289A	8	2	2	8			1	6															18			4	11	7	5									
G54E		2	2		1	8	7	5			1	15	7	1												2	5	5	4									
G54R		2			2	4	7				4	3	5	1		1		1							1			5	3								1	
G54V					4	1						4																	2	1								
G54W		2	2		2	2	7				3	5	2	2	1							1			1	1			1								6	4
G54E/I266N		4	2		1			2						5	1	1									2	5			1									1
G138C							2												3																			
G138S							1																															
F219I							1																															
F219S							1																															
M220I																																						
M220K						1		8						4	3	1													6	2								
M220R						9	2	2						8	3														3									
M220R														1	2														1	2								
M220T						3	1							1	2																							
M220V						8	5	1						1	12	1																						
I266N			1			1								2																								
G448S	2		1				1	2						1	2				1																			

<sup>a</sup>Values are cumulative numbers of isolates with the respective MIC values. Clinical breakpoints for results obtained with the CLSI methodology are not available; the values for S//R classification used here were adopted from reference 190.



the European Committee on Antimicrobial Susceptibility Testing (EUCAST) protocols were taken from previous references (16, 18, 24, 26, 38, 42, 44, 49, 54–81), and MIC values for tests done according to the Clinical and Laboratory Standards Institute (CLSI) protocols were taken from other previous references (3, 35, 43, 45, 50, 82–107).

Clinically used antifungal triazole compounds fall into two distinct structural classes (Fig. 1), which largely correlate with cross-resistance phenotypes: in *A. fumigatus* isolates with mutations in the *cyp51A* gene, cross-resistances within the substance pairs ITZ/PSZ and VRZ/ISAZ are most strongly correlated (16, 108). ISAZ and its azonium sulfate derivative for oral formulations have become an alternative for the therapy of invasive aspergillosis in patients with VRZ intolerance and seem to be more favorable than VRZ regarding drug-drug interactions (109).

In light of this, the use of at least VRZ along with ITZ as screening drugs is recommended to cover the range of currently known resistance phenotypes. Where available, extension of screening to the clinically relevant substances PSZ and even ISAZ may become advisable in the future.

### SPECIES IDENTIFICATION OF *ASPERGILLUS*

Exact species determination of molds by morphology, including that of *Aspergillus* spp., is a nontrivial issue (110). Molds can be slow to generate species-specific morphological traits allowing their discrimination, and they often require rather specific conditions to do so (111). On Sabouraud's agar plates, *Aspergillus* colonies appear as dense fields of conidiophores of various colors. Of the species discussed here, *A. flavus* appears yellowish to greenish, *A. fumigatus* and its close sibling species dark blue-green, *A. nidulans* green to dark yellow, *A. niger* black, and *A. terreus* light brownish (112). Due to the absence or delay of conidium formation, *A. fumigatus* may vary in its visual appearance to nearly white ("albino phenotype") (16, 111, 113). In addition, exact species boundaries are still not defined in all cases, and classification will ultimately require sequencing of several loci (e.g., internal transcribed spacer [ITS], *tub1*, *benA*, and *cmd* [114, 115]). Matrix-assisted laser desorption ionization–time of flight (MALDI-TOF) mass spectrometric species identification was recently extended to clinically important molds (116–118). In our own experience, commonly isolated species can readily be differentiated by using the commercial databases, and in cases where isolates with a common look (e.g., the bluish appearance of *A. fumigatus*) do not give rise to identification via MALDI-TOF, they turn out to be cryptic species not yet included in the scheme (119).

### SUSCEPTIBILITY TESTING PROCEDURES FOR *ASPERGILLUS* SPP.

There are several assays for susceptibility testing of *Aspergillus*, such as broth macrodilution (120), broth microdilution (121–123), and disk diffusion (124, 125) assays, Etest/MIC strip kits (126–128), and other commercial kits (129). Each method has distinct advantages and disadvantages; for example, macro- and microdilution assays are laborious and time-consuming, whereas Etest is easy to perform but expensive when used on a larger scale, and agar diffusion is inexpensive but not able to indicate exact MIC values (120).

Subsequent retesting and quantitative determination of MIC values by use of a reference method are also indicated to determine possible cross-resistance between compounds. In larger studies, two broth microdilution reference methods are mainly used for susceptibility testing: the CLSI M38-A2 protocol (121) and the EUCAST E.DEF 9.1 protocol (123, 130). These two standard methods are highly reproducible, and the MICs can be used for therapeutic selection of antifungal drugs as well as monitoring of antifungal drug susceptibility during treatment (120).

These two methods differ in several critical technical points, such as the amount of fungal inoculum ( $4 \times 10^4$  to  $5 \times 10^4$  CFU/ml as determined by use of a spectrophotometer for the CLSI method and  $2 \times 10^5$  to  $5 \times 10^5$  CFU/ml as determined by use of a hemocytometer for the EUCAST method). In addition, the EUCAST method uses a higher percentage of glucose (2%) in the test medium, which facilitates an increased

growth rate and specifies the use of flat-bottomed microtiter plates (120). It should be noted that both protocols have limitations in testing amphotericin B, which shows only a very narrow range of MIC values (0.5 to 2  $\mu\text{g/ml}$ ), as well as PSZ and the echinocandins (131). Because of their technical differences, the methods generate different absolute MIC values (131, 132), and consequently, divergent clinical breakpoints have been established by both EUCAST and CLSI (121, 133). EUCAST provides clinical breakpoints for antifungal agents for the interpretation of susceptibility testing results for fungi. Among molds, species-related clinical breakpoints have been determined only for *A. fumigatus*, *A. flavus*, *A. nidulans*, *A. niger*, and *A. terreus* (133). Using the EUCAST breakpoint table on these species, resistant isolates against the defined antifungal agents can be identified and subsequently investigated for mutations. There are no clinical breakpoints available from the CLSI, and therefore epidemiological cutoff values must be used (121).

In clinical studies, the results of *in vitro* susceptibility testing of *A. fumigatus* isolates did not necessarily correlate with the expected outcome of antifungal therapy in infected patients (123). From these studies, the “90-60 rule” was established, which implies that patients with infections caused by an *in vitro* susceptible strain will respond to the antifungal treatment in approximately 90% of cases, whereas patients suffering from infections caused by resistant isolates will respond in only about 60% of cases (120, 134). Therefore, although broth microdilution susceptibility screening is valuable for the selection of the best therapeutic agent, reliable prediction of the clinical outcome in a patient during antifungal therapy is still cumbersome (120, 130, 131). For these reasons, the combined use of molecular approaches to identify resistance and the phenotypic susceptibility testing described above may improve and guide successful treatment.


### EVOLUTION OF *cyp51* GENE FAMILIES

The major targets of azole antifungal drugs are cytochrome P450-type enzymes with sterol 14 $\alpha$ -demethylase activity. Inhibition of these enzymes leads to the formation of aberrant sterols, which do not complement the function of ergosterol in the plasma membrane and have cytotoxic and eventually fungistatic properties (135). The corresponding genes are called *cyp51* in molds and *ERG11* in ascomycetous yeast. P450 cytochromes are an evolutionarily ancient protein family and have undergone significant expansion and functional divergence across the entire phylogenetic tree. In molds, this is signified by at least three different lineages of paralogous genes, namely, *cyp51A*, *cyp51B*, and *cyp51C*.

Transcript and therefore protein abundance can be a direct function of gene copy number (“gene amplification” or “gene dosage effect”). If another copy of a *cyp51* gene is introduced, this can have a direct effect on drug resistance. For example, heterologous expression of a second *pdmA* copy in *A. nidulans* leads to strains that are selectable on azole-containing agar (136). This demonstrates that gene amplification may also contribute to tolerance, possibly as a prerequisite step to the expansion of *cyp51* gene families.

However, in evolutionary time spans, such duplicated genes will be fixed in a population only when they diverge in function (137). Paralogous genes may be lost and even may reemerge in populations (138).

Among *Aspergillus* spp., section Fumigati species contain *cyp51A* and *cyp51B* (139). Heterologous expression as well as biochemical testing has revealed differences between *A. fumigatus* Cyp51A and Cyp51B; despite identical substrate preferences and identical sensitivities against ITZ, VRZ, and PSZ, the isoenzymes showed differential tolerance of fluconazole, which was significantly increased for Cyp51A compared to that of Cyp51B (53, 140, 141). On comparing the Cyp51 protein sequences of molds to those of yeasts, e.g., Erg11 of *Candida albicans* (51), for which a multitude of substitutions have been brought forward in the context of azole resistance (e.g., see references 142 and 143), it becomes evident that their phylogenetic relationship is far from clear and that direct knowledge transfer to resistance in molds is difficult, at best. The proper



**FIG 4** Mutations in *A. fumigatus* *cyp51A* and its promoter region. (A) Alignment of the TR<sub>34</sub> (taken from isolate 168 [16]), TR<sub>46</sub> (taken from an environmental isolate [39]), and TR<sub>53</sub> (sequence taken from a Columbian isolate [41; P. LePape, personal communication]) promoter alleles with the wild-type sequences from *A. fumigatus* (*A. fu*), *A. fischeri* (*A. fish*), and *A. oerlinghausenensis* (*A. oerl*) (119). Green, TR<sub>34</sub> repeat unit; blue or red, 5' or 3' region and respective repeat sequences probably stemming from there. Black uppercase residues indicate sequence divergence in the repeat region, lowercase residues indicate adjacent residues, and black boxes indicate TR<sub>34</sub>, TR<sub>46</sub>, and TR<sub>53</sub> repeats. (B) In sequencing data, these three different types of *A. fumigatus* can easily be differentiated by the 5' upstream sequence preceding the static repeat unit: sequences different from the wild-type sequence are indicated by underscored nucleotides. (C) Known amino acid substitutions in Cyp51A. Red, known resistance-conferring substitutions; black, substitutions present in the population, probably without an effect on drug susceptibility. (D) *cyp51A* gene features visualized by use of sequence analysis software (Geneious R10). The data are available as a .gb annotated file or as a Word document in a community-editable form from [https://github.com/oliverbader/Aspergillus\\_fumigatus\\_cyp51A](https://github.com/oliverbader/Aspergillus_fumigatus_cyp51A).

paralog to *C. albicans* *ERG11* is probably *cyp51B* (29). To complicate matters, due to differences in expression patterns (Cyp51B is expressed constitutively, and Cyp51A is inducible), Cyp51A is believed to be the major player in azole resistance in *A. fumigatus*.

Among section Flavi species, at least for *A. flavus* and *A. calidoustus*, the additional paralog *cyp51C* is present, and in *A. calidoustus* *cyp51A* has been lost (144). In these species, Cyp51C appears to be the key player (see below).

## RESISTANCE THROUGH TRANSCRIPTIONAL MODULATION OF *cyp51* GENES

To date, three different azole resistance-associated variations of the *cyp51A* promoter (Fig. 4A) are known for *A. fumigatus*. Most frequently, two tandem repeats, of 34 and 46 bp, have been described to occur 279 bp upstream of the coding region (84, 98, 145). The TR<sub>34</sub> promoter has been found only with the L98H exchange conferring high ITZ resistance, and the TR<sub>46</sub> promoter has been found in conjunction with the Y121F and T289A exchanges in intermediate to highly VRZ-resistant isolates. Additionally, a 53-bp repeat has been found in isolates that are cross resistant to VRZ and ITZ, without

any further Cyp51A alterations (41, 145). Most recently, isolates with up to three or four copies of the TR<sub>46</sub> repeat were identified (81). Changes in the promoter region must be interpreted in the context of gene expression levels (Fig. 2D). For the TR<sub>34</sub> promoter, a 2-fold higher activity in a reporter assay (146) and an 8-fold higher *cyp51A* gene expression level (84) have been reported. However, this remains untested so far for any other promoter variants. Molecular studies of the altered *A. fumigatus cyp51A* promoter have identified opposing actions of the CCAAT binding complex (CBC), which is involved in the regulation of many genes in eukaryotes by binding CCAAT sites (19, 147, 148), and the HAP complex, an element of the sterol response pathway (20). Genome sequencing of matched resistant and susceptible isolates also identified a mutation in *hapE*, introducing the P88L amino acid exchange, which was shown to be responsible for the azole resistance phenotype (19). Detailed molecular studies of the HAP complex have found it to interact with the *cyp51A* promoter at the repeat elements found in drug-resistant isolates, there opposing the action of the sterol response factor SrbA (20). The P88L substitution leads to a significant loss of binding activity, in turn leading to unopposed action of the sterol biosynthesis response pathway, upregulating *cyp51A*, among other sterol biosynthesis genes. As it has been reported only once, the number of clinical isolates containing the P88L substitution in HapE appears not to be very high, or at least underinvestigated.

Outside *A. fumigatus*, data on *cyp51* promoter alterations are scarce. Only in *Aspergillus (Neosartoriya) fischeri* and *Aspergillus oerlinghausense*, the two known species most closely related to *A. fumigatus*, can a similar to identical 34-bp core repeat region readily be identified in the respective promoters of the corresponding *cyp51A* genes (Fig. 4A). In *A. oerlinghausense*, a short insertion upstream of the core repeat correlates with the decreased azole susceptibility in the only two existing isolates of this new species (119), albeit molecular confirmation for this observation is currently lacking. The 1-kb regions flanking the three *cyp51* paralogs of resistant and susceptible *A. flavus* isolates investigated (149) did not deviate from each other, but further upstream a short 4-bp deletion was found in two resistant strains (149). The role of this deletion remains unclear. In sequencing data for *A. fumigatus cyp51A*, the type of repeat can easily be identified by the first 10 bp preceding the static 34-bp repeat unit (Fig. 4B).

### RESISTANCE-CONFERRING ALTERATIONS WITHIN Cyp51 PROTEINS

Mair et al. (29) recently suggested the use of a positional nomenclature for fungal resistance mutations to allow easier transfer of knowledge on orthologous proteins of different species. Based on a preformed alignment of the amino acid sequences, the respective number in a reference is given instead of the residue number of the affected amino acid itself. For Cyp51A-orthologous proteins, the reference was suggested to be Cyp51A of *A. fumigatus*, and for Cyp51B-orthologous proteins, the reference is Cyp51B of *Zymoseptoria tritici* (anamorph of *Mycosphaerella graminicola*). For Cyp51C, no suggestion has yet been made; for the purpose of this review, we therefore name Cyp51C of *A. flavus* as the reference. The positional nomenclature is used below and summarized in Table 3.

#### *Aspergillus fumigatus*

In approximately 50 to 80% of the clinically observed ARAf isolates, resistance is attributable to amino acid exchanges within the Cyp51A protein (16–18, 95). Cyp51B substitutions seen in the population apparently do not contribute to resistance. Similarly, only a subset of Cyp51A substitutions present in the population lead to resistance against azoles (Fig. 4C). Notably, several synonymous (silent) mutations in *cyp51A* are also known, e.g., at the L358 or C454 codon (82) or the L77, R65, or E356 codon (150). These are interesting in a phylogenetic context but have no relevance for deduction of resistance phenotypes. On encountering nonsynonymous mutations in sequencing data, these must be discriminated from allelic variants without any effects on azole susceptibility. Several substitutions (e.g., F46Y, H147Y, M172V, N248T, N248K, D255E, D343N, E427K, and G434C) seem not to be involved in azole resistance, as they are also

**TABLE 3** Cyp51 substitutions in non-*A. fumigatus* organisms and their correspondence to the reference positions used in the text

Organism	Reference position <sup>a</sup>			Observed substitution		
	Gene	Position <sup>b</sup>	Reference	Gene	Substitution	Reference
<i>A. flavus</i>	<i>Afcyp51A</i>	Y68	29	<i>cyp51A</i>	Y132N	160
	<i>Afcyp51A</i>	K133	29	<i>cyp51A</i>	K197N	160
	<i>Afcyp51A</i>	NA		<i>cyp51A</i>	A205T	149
	<i>Afcyp51A</i>	D280	29	<i>cyp51A</i>	D282E	160
	<i>Afcyp51A</i>	M286	29	<i>cyp51A</i>	M288L	160
	<i>Afcyp51A</i>	T470	29	<i>cyp51A</i>	T469S	160
	<i>Ztcyp51B</i>	H430	29	<i>cyp51B</i>	H399P	160
	<i>Ztcyp51B</i>	A453	29	<i>cyp51B</i>	D411N	160
	<i>Ztcyp51B</i>	T496	29	<i>cyp51B</i>	T454P	160
	<i>Ztcyp51B</i>	NA		<i>cyp51B</i>	T486P	160
	<i>Afcyp51C</i>	M54	Proposed here	<i>cyp51C</i>	M54T	149
	<i>Afcyp51C</i>	S196	Proposed here	<i>cyp51C</i>	S196F	161
	<i>Afcyp51C</i>	S240	Proposed here	<i>cyp51C</i>	S240A	149, 162
	<i>Afcyp51C</i>	D254	Proposed here	<i>cyp51C</i>	D254N	149
	<i>Afcyp51C</i>	I285	Proposed here	<i>cyp51C</i>	I285V	149
	<i>Afcyp51C</i>	Y319	Proposed here	<i>cyp51C</i>	Y319H	149, 191
	<i>Afcyp51C</i>	A324	Proposed here	<i>cyp51C</i>	A324P	161
	<i>Afcyp51C</i>	N423	Proposed here	<i>cyp51C</i>	N423D	161
	<i>Afcyp51C</i>	V465	Proposed here	<i>cyp51C</i>	V465 M	161
<i>A. terreus</i>	<i>Afcyp51A</i>	M220	29	<i>cyp51A</i>	M217I	165
<i>A. niger</i>	<i>Afcyp51A</i>	K230	29	<i>cyp51A</i>	R228Q	166
<i>A. tubingensis</i>	<i>Afcyp51A</i>	L21	29	<i>cyp51A</i>	L21F	166

<sup>a</sup>Accession numbers for reference sequences are as follows: *A. fumigatus* *cyp51A*, [AF338659](#); *Zygomycetia tritici* (anamorph of *Mycosphaerella graminicola*) *cyp51B*, [AY253234](#); and *cyp51C*, [AKQ20794.1](#).

<sup>b</sup>NA, not applicable (the respective residue was not found at this position in publicly deposited *A. flavus* sequences).

found in different combinations in azole-susceptible isolates (56, 62, 70, 75, 82, 87, 151), and these sites are mostly localized at the protein surface (152). Among these, isolates harboring the multiply substituted *cyp51A* F46Y/G89G/M172V/L358L/E427K/C454C allele (18, 56) appear to form a separate phylogenetic subgroup rather than grouping with strains with acquired resistance. Nevertheless, these substitutions do occur alongside those with relevance for reduced antifungal drug susceptibility, e.g., the TR<sub>34</sub>/L98H substitution (Fig. 4 and Table 2), likely where the original resistant strain crossed with another isolate of the population (153, 154).

The Cyp51A-I266N substitution reported from Japan (90) can probably be discarded as an interpretation error, as the reference sequence already has an asparagine (N) residue at this codon and it was not observed in subsequent studies (155).

Azole-based inhibitors of Cyp51 interact with the protein by complexing the heme group located in the catalytic center (152). For amino acid substitutions located along the periphery of the protein (e.g., S52T, Q88H, N125I, Q141H, S297T, or F495H), homology modeling suggests that they do not relevantly change its functionality (82, 152). Consequently, residues where substitutions lead to azole resistance, e.g., G54 (57, 83, 90, 103), G138 (56, 75, 104), or M220 (56, 82, 99), are mainly located close to the opening of one of the two ligand access channels of the protein, blocking the docking of most azole molecules (82, 152). Similarly, the L98 residue is located on a highly conserved, loop-forming, arch-like structure affecting the entry to the ligand access channel (82).

For several of the polymorphisms seen in antifungal drug-resistant isolates, the resistance-conferring nature of the particular substitution has been confirmed on a molecular level: biochemical testing of purified recombinant *A. fumigatus* Cyp51A-M220K and Cyp51A-G54W expressed in *Escherichia coli* (141) showed inhibition kinetics that would be expected from the respective clinical isolates' phenotypes, namely, a strong reduction in inhibition by ITZ and PSZ but not by VRZ. Interestingly, the G54W mutation does occur in both isolates with increased and those with susceptible MIC values (Tables 1 and 2). This raises the question of whether this substitution is actually relevant to clinical resistance, despite the fact that its contribution to reduced suscep-



tibility has been confirmed *in vitro* (141). This may also be an example of interactions with other factors, such as protein-protein interactions potentially dependent on the genetic strain background.

A very different situation was found for Cyp51A-L98H. In this case, inhibition of resistance was only marginally impaired, but the enzyme retained some residual activity at higher triazole concentrations, thus likely explaining its *in vivo* resistance phenotype (141). Structural modeling suggests that the L98H substitution is located at neither the substrate binding cleft nor the central heme group, but rather in a conserved surface arch, which would support these data (152).

The gold standard for *in vivo* testing of this is heterologous expression of a potential resistance-conferring allele in a susceptible host. In the case of *cyp51A* mutations from *A. fumigatus*, susceptible strains of the fungal model organism *Saccharomyces cerevisiae* and, more recently, *A. fumigatus* itself have been used.

*S. cerevisiae* is readily available for genetic studies and has also served as a host in several studies of azole resistance in several plant-pathogenic fungi (11). *S. cerevisiae* encodes only one Cyp51 ortholog, namely, Erg11. However, deletion of *ERG11* results in lethality, which is why complementation experiments were done in an *ERG11* tetracycline-repressible strain (140) or by a one-step gene replacement strategy (75). In these studies, introduction of the *cyp51A* or *cyp5B* gene was shown to complement the loss of the respective *S. cerevisiae* enzyme function.

Using a tetracycline-regulatable system, Alcazar-Fuoli et al. (156) were able to confirm the anticipated phenotypes for several mutations at positions G54 and M220: elevated ITZ MICs were found for G54E, G54V, G54W, and M220K mutants, and the M220K and M220I substitutions conferred VRZ resistance. The highest MIC values for PSZ were found for the G54W mutant, followed by the G54V, G54E, and M220K mutants. Using the gene replacement strategy, Albarrag et al. (75) confirmed decreased azole susceptibility for Cyp51A-G138C and Cyp51A-Y431C. Unexpectedly, the G434C mutant showed no resistance phenotype. The authors suggested that this method may not be suitable for alterations at the far C-terminal end of the protein, where interactions with as yet unknown binding partners may be the reason for reduced susceptibility in the original host.

Although a high degree of cross- and pan-azole resistance is seen, the quantitative degree of reduced drug sensitivity still varies between individual substitutions as well as compounds (Table 2). These effects may also be cumulative in cases where different promoter modifications and/or amino acid substitutions are combined in a single isolate. For *A. fumigatus*, these effects have been best investigated for the TR<sub>46</sub>/Y121F/T289A allele. Isolates with this allele show high VRZ MICs along with reduced ITZ and PSZ susceptibilities (45). A single isolate with only the Y121F substitution showed only an intermediate VRZ MIC (157). Based on this, molecular dissection of this allele in an *A. fumigatus* expression system showed that the combination of TR<sub>46</sub> with the Y121F mutation was sufficient to induce both high ITZ and VRZ resistances, while TR<sub>46</sub> or the T289A mutation alone or in combination did not have a similar effect. Most strikingly, the strain arising from combination of the T289A mutation with the TR<sub>46</sub>/Y121F allele was again less resistant to ITZ (158). Others have also found the TR<sub>46</sub>/Y121F/T289A allele in combination with the G448S substitution in an isolate with pan-azole resistance (67). However, the G448S amino acid change can result in cross-resistance to VRZ and ITZ (159). The MIC values reported for the individual mutations are summarized in Tables 1 and 2, stratified by testing method.

### ***Aspergillus flavus***

Next to *A. fumigatus*, most other amino acid substitutions in Cyp51 proteins have been described for *Aspergillus flavus*. A key difference of this species from other *Aspergillus* species is the existence of a third paralog in *A. flavus*, namely, *cyp51C*. Substitutions have been observed in all three paralogs. Using the positional nomenclature (Table 3), these were at Y68, K133, D280, M286, and T470 in Cyp51A (149, 160),



	54								431					
<i>A. fumigatus</i> <b>cyp51A</b>	TISY <b>G</b> IDPYK	QEVRSKLTA	ARMRSIYVDIINQRRLDGDKDSQ	EIAHMMITL	LEELYQEQLANLGP	TKEQENDE-VVD <b>Y</b> GYG	VNLGVILAT							
<i>A. flavus</i> NRRL3357	TIHYG <b>M</b> DPYG	EEVRSKLTT	ARMRSIYIDIINKRRNAGDNVPE	EIAH <b>I</b> MITL	VEEL <b>Y</b> QEQLANLER	PQEN <b>N</b> KD-DIVDYG <b>Y</b> G	LNLA <b>V</b> IVAT							
<i>A. flavus</i> AF70	..... <b>T</b> .....	..... <b>P</b> .....	..... <b>A</b> ..... <b>N</b> .....	..... <b>I</b> .....	..... <b>Y</b> ..... <b>A</b> .....	..... <b>D</b> .....	..... <b>V</b> .....							
	*	*	*	*	*	*	*							
	54	196	240	254	285	319	323	423					465	
<i>A. oryzae</i> RIB40	.....	.....	....A.....N....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
<i>A. nomius</i>	.....	.....	....A.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
<i>A. lentulus</i>	..S..I...K	Q.....A	.....V...Q...LD.....	.....M....	L.....	AK.QEN.-EV.....	V..IG..L..							
<i>T. interdigitale</i>	..S.....K	.....	....A..T...E...RE.GKDSQ	.....M....	.....R.....GP	.NP.E.EE.M.....	V..T.....							
<i>T. tonsurans</i>	..S.....K	.....	....A..T...E...RE.AKRNH	.....M....	.....R.....PH	.NP.E.EE.M.....	V..T.....							
<i>T. rubrum</i>	..S.....K	.....	S...A..TE...E...RE.AKRNH	.....M....	L...R.....PH	.NTDEQEE.M.N....	V..T.....							
<i>T. soudanense</i>	..S.....K	.....	S...A..TE...E...RE.AKRNH	.....M....	L...R.....PH	.NTDEQEE.M.N....	V..M....T							
<i>M. canis</i>	..R..L...R	.....	....A..T...E...R..T.INA	.....	...F.....P.	.K.DE.E-.M.....	V..T...T							
<i>H. capsulatum</i>	..R.....K	.....A	TL.HA..L...E...R.AR.ANG	...N...S.	..Q.....D..P.	.KVDE.E-.M.....	V..T...I							
<i>B. dermatitidis</i>	..C..I...K	.....	...HA.....D...LA.TTSDA	...N...S.	..K.....S..PQ	.KLEE.E-.M.....	...T...T							

**FIG 5** Amino acid substitutions in Cyp51C proteins. Numbers and asterisks show the reference positions of residues discussed in the text (blue). Vertical dashes, sequence identity between *A. flavus* NRRL3357 Cyp51C (accession number [XP\\_002383931](#)) and *A. fumigatus* Cyp51A (accession number [AF338659](#)); dots, residues identical to those in *A. flavus* NRRL3357 Cyp51C. Accession numbers for the other reference sequences are as follows: *A. flavus* ATCC MYA-384/AF70, [KOC15064](#); *A. oryzae* RIB40, [AB514682](#); *A. nomius*, [XP\\_015411243](#); *A. lentulus*, [AEB77687](#); *Trichophyton interdigitale*, [EZ31978](#); *Trichophyton tonsurans*, [EGD95049](#); *Trichophyton rubrum*, [XP\\_003235929](#); *Trichophyton soudanense*, [EZ372647](#); *Microsporum canis*, [XP\\_002845046](#); *Histoplasma capsulatum*, [EER42982](#); and *Blastomyces dermatitidis*, [EGE84227](#).

H430, D435, and T496 in Cyp51B (160), and M54, S196, S240, D254, Y319, A324, N423, and V465 in Cyp51C (149, 161, 162).

As in *A. fumigatus* Cyp51A, the relevance of the Cyp51 substitutions seen in *A. flavus* must also be interpreted with respect to the variability of the respective positions in the population as well as in other fungal species.

Both the M54T and S240A substitutions in Cyp51C are present in susceptible isolates and were identified as variants in the respective studies (149, 161) only because the genome-sequenced strain MYA-384/AF70 (accession number [PRJNA217045](#)) contains a *cyp51C* isoform divergent from that of the general *A. flavus* population, including the reference genome (*A. flavus* NRRL3357 [accession number [PRJNA13284](#)]) (163). The latter also applies to the S196P, D254N, and N423D substitutions (Fig. 5). Only in the case of the M54T substitution is a known resistance substitution of Cyp51A (G54) located in the close vicinity; however, Cyp51A G54 actually corresponds to the adjacent Cyp51C position G53, not M54 (Fig. 5). At all positions except S196, variation is seen between other species. The S240A substitution (note that due to the intron in *A. flavus cyp51C*, the respective mutation is referred to as a T788G substitution; nucleotide positions of the mutations are listed in the work of Paul et al. [149]) remains particularly controversial. In one study, it was associated with a high VRZ MIC, and this phenotype was confirmed using molecular complementation assays (162). However, others have also observed it in susceptible isolates (149). Together, these observations make involvement of these residues in drug resistance unlikely (149).

Of the four remaining residues (I285, Y319, A232, and V465), only I285 shows variation within the *Aspergillus* genus (161), but the reported I285V substitution co-occurred only in isolates with other potential resistance-conferring substitutions (149).

For *cyp51A* and *cyp51B*, microevolution experiments using long-term incubation of *A. flavus* on azole-containing media have yielded strains with reduced azole susceptibility and mutations leading to multiple amino acid exchanges in both Cyp51A and Cyp51B (160). This was most prominent for Cyp51A-K197N (position K133), which was observed three times independently. Other substitutions (Table 3) were observed only once, and several less susceptible isolates with unaltered *cyp51A* and *cyp51B* sequences were also observed. Potential changes in *cyp51C* or other genes were not investigated, limiting these observations; similarly, the K133 position is not known outside these experiments to hold resistance substitutions and has not been verified independently.

**TABLE 4** PCR-based assays to characterize molecular resistance azole mechanisms in *Aspergillus* spp.

PCR use and format	Target	Reference(s)
Analyses of <i>A. fumigatus</i> <i>cyp51A</i>		
Real-time PCR	Single regions (~100 bp)	192, 193
Mixed-format real-time PCR	TR- and amino acid substitution-specific regions	92, 194
PCR-RFLP	Multiple single regions (180–270 bp)	167
Nested PCR, two amplicons	F1, TR region to L98; F2, G54 to N266	193
Conventional one-step and nested PCR <sup>a</sup>	Single regions (100–173 bp) for TR <sub>34</sub> /L98H and M220	175
	Single regions (103–143 bp) for TR <sub>46</sub> and the L98H substitution	174
PCR with TaqMan probes	TR, G54, L98, G138, M220	83
	L98H and Y121F substitutions	71
Primer extension	TR <sub>34</sub> and 15 polymorphic sites	151
LightCycler endpoint genotyping	L98H, Y121F substitutions	71
AsperGenius (PathoNostics BV, Maastricht, The Netherlands)	TR <sub>34</sub> , L98H, Y121F, T289A substitutions	25, 107, 176,
multiplex real-time PCR and melting curve analysis <sup>a</sup>		177
Real-time RT-PCR	<i>cyp51A</i> gene expression	75
Analyses of <i>A. fumigatus</i> genes other than <i>cyp51A</i>		
PCR plus sequencing	Full <i>cyp51B</i> gene in four amplicons	75
Real-time RT-PCR	<i>cyp51B</i> gene expression	75
	<i>mdr1</i> and <i>mdr4</i> gene expression	75
PCR with TaqMan probes	<i>cyp51B</i> S35, P294	151
PCR plus sequencing	Full <i>hapE</i> gene	19
PCR plus sequencing, real-time RT-PCR	<i>cyp51A</i> , <i>mdr1-4</i> , <i>atrF</i> , <i>cdr1B</i> ( <i>abcG1</i> ), <i>mfs56</i>	17
Analyses of <i>A. flavus</i> <i>cyp51</i> genes		
PCR plus sequencing	<i>cyp51A</i> and <i>-B</i> genes plus 5' and 3' regulatory sequences	160
PCR plus sequencing, TaqMan probes	<i>cyp51A</i> , <i>-B</i> , and <i>-C</i> genes plus 5' and 3' regulatory sequences	162
PCR plus sequencing	<i>cyp51A</i> , <i>-B</i> , and <i>-C</i> genes plus 5' and 3' regulatory sequences	149

<sup>a</sup>Non-culture-based, direct investigation of clinical (respiratory) samples was possible after confirmation of *Aspergillus* DNA in the sample by use of diagnostic *Aspergillus* PCR assays.

How any of these substitutions affect antifungal therapy in a clinical setting remains unclear. In a nonimmunocompromised mouse model, VRZ treatment was not relevantly impaired by mutations in *cyp51C* (164). The authors of that study discussed the position of the Y319H mutation as being rather far from the binding position of azoles and a potential loss of fitness in resistant strains as explanations for the unexpected results.


### Other *Aspergillus* Species

Data on azole resistance and the underlying mutations in species apart from *A. fumigatus* and *A. flavus* are scarce; however, there are 3 single observations of Cyp51A alterations reported in the literature (Table 3): the M217I substitution (position M220) in *A. terreus* (165), the R228Q substitution (position K230) in *A. niger* (166), and the L21F substitution (position L21) in *A. tubingensis* (166). The isolate with the R228Q substitution showed a PSZ MIC above the epidemiological cutoff value. For the *A. tubingensis* isolate with the L21F substitution, especially the measured ITZ MIC value of 64 µg/ml was highly elevated compared to those for other isolates (166). Only the M217I substitution corresponds to a well-known residue (M220) involved in azole resistance in *A. fumigatus*. Neither K230 nor L21 has been implicated in Cyp51A-based drug resistance, and neither observed exchange has yet been validated independently.

### MOLECULAR METHODS FOR DETECTION OF ALTERATIONS AT *cyp51* LOCI

#### *Aspergillus fumigatus*

A standard method to detect alterations of the *cyp51A* gene in fungal isolates is not yet defined. The most commonly used method is simple PCR amplification of the entire coding and promoter region, followed by Sanger sequencing of the PCR products. Identification of the promoter repeat type can easily be achieved by analyzing the 5' upstream sequence of the static repeat unit (Fig. 4B). Specific methods for detection of the TR<sub>34</sub> and TR<sub>46</sub> variants are also available, including several rapid TaqMan-based approaches and melting curve analyses on LightCycler instruments (Table 4). Utilization of an AluI-based restriction fragment length polymorphism (RFLP) method (167) at the



**FIG 6** Interpretation of AluI-digested *A. fumigatus cyp51A* PCR product surrounding the L98 position. (A) In isolates carrying the L98H substitution, the AluI restriction site at this codon is abolished (167). (B) A PCR product amplified with the P5-P7 primer pair (16) encompasses both the L98 site and the promoter repeat region. When digested with AluI, the wild-type product is cleaved into four fragments, and the length of fragment A is characteristic of the length of the TR region. In L98H isolates, fragments B and C are not cleaved apart, resulting in a visible size shift. (C) When resolved in a high-percentage (e.g., 2.5%) agarose gel, the size shift is visualized as specific RFLP patterns for TR<sub>34</sub>/L98H- and TR<sub>46</sub>-carrying isolates. Other substitutions cannot be found using this method, as their RFLP patterns correspond to the wild-type patterns. The method may, however, be suitable to detect other promoter variants.

L98 codon probably has the highest value for most laboratories, as the assay is readily performed without the need for specialized equipment. It allows easy simultaneous detection of both TR<sub>34</sub> and TR<sub>46</sub> repeats (and likely also others), as well as the L98H substitution (Fig. 6), therefore encompassing the most frequently encountered resistance-associated *cyp51A* alleles. The commercially available AsperGenius assay also covers these two alleles (Table 4). For a few selected other exchanges at polymorphic sites, namely, G54 and M220, specialized PCR-based detection methods have been developed (Table 4).

For all other mutations, including so far unknown ones, PCR amplification of the gene in either a single (3, 18, 45, 50, 68, 82, 104, 168) or multiple (16, 44, 54, 56, 60, 62, 63, 75, 84, 100, 157, 169–171) amplicons plus subsequent sequencing is necessary. Identification of mutations in trace files or derived fasta sequences and of the respective amino acid substitutions is most easily achieved by mapping against a reference sequence. To facilitate this, we created an annotated reference containing currently known polymorphic sites along with other relevant features (Fig. 4D) ([https://github.com/oliverbader/Aspergillus\\_fumigatus\\_cyp51A](https://github.com/oliverbader/Aspergillus_fumigatus_cyp51A)). Alternatively, there is also an online tool available to perform sequence analysis on *A. fumigatus cyp51A* (172).

Given the low pathogen yields by cultural methods for patients with hematological malignancies who are at high risk for invasive aspergillosis, it is very likely that azole resistance is still underdiagnosed in this population (173). The reported high mortality rates associated with ARAF underline the necessity of using non-culture-based assays for the detection of both *Aspergillus* spp. and azole resistance directly from clinical samples (24, 25, 174, 175). This approach requires higher sensitivity and lower detection thresholds than those for working on cultured isolates. Molecular methods for detecting *cyp51A* alterations directly from clinical samples have to combine high sensitivity and specificity to ensure the amplification of small amounts of *Aspergillus* DNA and to avoid cross-reactivity with human DNA. Most formats are PCR-based methods with subsequent DNA sequence analysis for the detection of mutations. PCR assays investigating primary clinical samples directly do not have the ability to amplify the whole *cyp51A* gene due to the very small amount of intact fungal high-molecular-weight DNA

in these samples. Therefore, individual PCR assays were established to amplify the key mutation-carrying gene sections. In this scenario, nested PCR assays using two consecutive PCRs amplifying the same gene region have shown the highest sensitivities. A new real-time PCR-based system for detection of *Aspergillus* DNA and the TR<sub>34</sub>/L98H and Y121F/T289A *cyp51A* alterations without subsequent DNA sequencing is the AsperGenius system from PathoNostics (25, 107, 176, 177). This test system is based on *Aspergillus* DNA amplification by multiplex real-time PCR followed by melting curve analysis to detect the mutations in different PCR cyclers-based detection channels. Protocols that have been used successfully directly on clinical samples (bronchoalveolar lavage [BAL] fluid and blood samples) with low pathogen DNA amounts, e.g., nested PCR (174) or multiplex PCR (25, 58, 176, 177), are summarized in Table 4. In this rapidly advancing field, prospective studies are still ongoing.

### Other *Aspergillus* Species

For *Aspergillus* spp. other than *A. fumigatus*, there are currently no commonly used protocols and—to date—no commercially available kits. The best-characterized species is *Aspergillus flavus*. Molecular analysis of the *cyp51A* and *cyp51B* genes for resistance mutations was performed using PCR with subsequent DNA sequence analysis (160). In another study, amplification of *cyp51A*, *-B*, and *-C* included an additional 1 kb of genomic DNA up- and downstream of the coding regions (162). Assays for the analysis of *cyp51* expression levels by quantitative TaqMan-based real-time PCR have also been described (162). The aspects of molecular azole resistance detection in *A. flavus* are also summarized in Table 4.

### OTHER RESISTANCE MECHANISMS

For a significant proportion of azole-resistant clinical *A. fumigatus* isolates, no mutations within the *cyp51A* locus are observed (16–18, 95). Compilation of the MIC values from the literature (Tables 1 and 2) showed that isolates with the VRZ resistance-conferring TR<sub>46</sub>/Y121F/T289A allele fall into two clearly distinct groups with respect to ITZ cross-resistance, namely, VRZ<sup>r</sup> ITZ<sup>s/i</sup> and VRZ<sup>r</sup> ITZ<sup>r</sup>, in the absence of other *cyp51A* mutations. We also previously observed an unusual TR<sub>34</sub>/L98H isolate with an additional VRZ MIC of >32 µg/ml (39). Together, these data highlight the multifactorial ability of the fungus to overcome susceptibility, so the genetic basis for resistance is not restricted just to mutations within the *cyp51A* region. However, so far, only a few other mechanisms have been discussed to contribute to clinical azole resistance.

Most importantly, the expression of ABC transporters can be upregulated in clinical isolates (Fig. 2E); however, their contribution to azole resistance has not yet been fully clarified (17, 23). So far, the best-studied transporter in correlation with drug resistance is the ABC efflux transporter Cdr1B. This transporter shows high homology to the efflux pump Cdr1p from *Candida albicans* (CaCdr1), which is regulated by the transcription factor Tac1. Overexpression of CaCdr1 increases the resistance to fluconazole in both *A. fumigatus* (17, 21) and *C. albicans* (178). A potential regulator of Cdr1B in *A. fumigatus* that is similar to Tac1 in *C. albicans* is the zinc cluster transcription factor AtrR (179). Mutations in this regulator might provide explanations for increased resistance mediated by overexpression of Cdr1B. Nevertheless, gain-of-function mutations of either the *atrR* or *cdr1B* coding region or their promoters have not been described so far.

Data from *in vitro* approaches, such as mutagenesis, genetic backcrosses, and complementation in *S. cerevisiae*, have led to the identification of other potential drug resistance genes (19, 22, 180–182), but these have not yet been confirmed to exist in clinical or environmental isolates.

Especially when interpreting PCR and/or sequencing data in the absence of a culture isolate for susceptibility testing, it must be kept in mind that the presence of a particular mutation likely indicates resistance toward specific compounds (high positive predictive value); however, the absence of a mutation per se does not indicate susceptibility (low negative predictive value).

Standardized phenotypic tests that can be used for diagnostic procedures for detection of increased sterol biosynthesis or efflux have not yet been proposed.

### CONTRIBUTION OF PHYLOGENETIC ANALYSES TO RESISTANCE DETECTION

Using genetic strain typing of a small *A. fumigatus* strain set, it has been suggested that the TR<sub>34</sub>/L98H allele arose only once and has since been propagating through the *A. fumigatus* population (35). For isolates with the TR<sub>46</sub>/Y121F/T289A or G54E/W allele, similar observations have been made (39, 108). This has led to only a few distinct lineages, which can be differentiated by both short-tandem-repeat typing (e.g., see references 24 and 108) and *csp1* typing (e.g., see references 39 and 170). Because of this and a significant degree of cross-resistance to agricultural fungicides, the hypothesis of an environmental origin of these alleles has been suggested (50). Strains with these alleles are thought to be propagated in the environment through the use of different azole compounds by the farming industry. Along with that proposition, an exogenous route of infection of patients has been proposed for these isolates. In contrast, isolates with other exchanges at G54 or at M220 or with one of the rarer mutations are more likely to have originated within the patient under prolonged antifungal therapy with mold-active azoles.

Short-tandem-repeat typing of *A. fumigatus* (STRAf typing) utilizes the copy number variability of nine short repeat sequences, of 2 to 4 bp in length, whose PCR amplicons are resolved by capillary electrophoresis (183). STRAf typing has a very high discriminatory power (183), and for ARAf isolates, it has been shown to readily demonstrate the genetic nearness of globally isolated TR<sub>34</sub>/L98H or G54E strains (108). This makes it an excellent and robust reference method; however, capillary electrophoresis requires special equipment, which rules out its use as a rapidly established laboratory technique. Thus, the typing approach using the repetitive region of the *csp1* gene is implemented more easily (184–186). The *csp1* gene is simply amplified by PCR and sequenced, and the sequence is searched for the number and type of 12-bp-long elements. Their order gives one of (currently) 27 *csp1* types (184). As a single-locus method, it has a lower discriminatory power than that of STRAf typing, but it is more easily implemented and is practical even for a single isolate. Among the isolates tested in our own laboratory, there was a moderate correlation of TR<sub>34</sub>/L98H isolates with the t04B *csp1* type, a type which has not been described for isolates with any other *cyp51A* allele (39).

Additionally, sequencing of hypervariable regions in cell wall genes (“TRESP”) (187) has been shown to yield a sufficiently diverse marker set to discriminate within the *A. fumigatus* population. By this analysis, TR<sub>34</sub>/L98H strains also formed a less diverse (i.e., more closely related) subpopulation.

Ultimately, whole-genome sequencing will have to reveal the phylogenetic and evolutionary relationships among early TR<sub>34</sub>/L98H *A. fumigatus* isolates and their progeny. A first study (154) did indeed find similar rates of recombination between strains isolated more distantly as well as in close proximity. Similar conclusions of both local clonal expansion and global recombination were drawn from a study using STRAf typing (188).

From a clinical point of view, phylogenetic analyses are therefore interesting for identifying local outbreaks or common sources of infection, but they unfortunately have—despite a likely common ancestral origin for strains with the major resistance alleles—only low value for the prediction of azole resistance phenotypes in *A. fumigatus*.

### CONCLUSIONS

The emergence of *Aspergillus* strains with azole resistance, and even pan-azole resistance, spreading through the environment rather than originating under therapy in individual patients poses a significant threat to vulnerable patient groups. Although this has not yet been shown for other alleles, most worrying is that at least Cyp51A-G54W- or Cyp51A-M220K-carrying azole-resistant *A. fumigatus* strains do not seem to suffer from growth or fitness defects (189): as in phytopathogens, azole resistance in



human-pathogenic *Aspergillus* spp. is likely here to stay. Recombination with other alleles has already been observed within *A. fumigatus* *cyp51A* alleles, as both TR<sub>34</sub>/L98H and TR<sub>46</sub>/Y121F/T289A alleles have been observed in combination with substitutions seen in the population (39, 81, 154). Since *A. fumigatus* does have a sexual cycle and genetically crosses in the wild (153, 154), it stands to reason that despite the initial clonal expansion, phylogenetic relationships between the different loci and resistance alleles will blur and eventually dissipate in the future. Within a species, the resistance traits may freely combine with each other as well as with other traits present in the wild, leading to increasingly resistant populations. Establishing the local epidemiology, for which we outlined potential workflows here, and setup of proper patient testing procedures are therefore indicated in clinical settings.

## ACKNOWLEDGMENTS

This work was supported by a research grant from Pfizer Germany to O.B. (grant W1186302) and by a research grant of Burapha University, through the National Research Council of Thailand, to M.T. (grant 165/2559).

We thank Agnieszka Goretzki (University Medical Center Göttingen, Göttingen, Germany) for conducting the RFLP analysis in Fig. 6. We thank Patrice Le Pape (Université de Nantes, Nantes, France) for providing the unpublished sequence of the 53-bp promoter repeat unit for Fig. 4. The anonymous reviewers are thanked for pointing out several gaps in the initial manuscript.

All authors contributed to writing and approved the final version of the manuscript.

D.B. serves in the speakers' bureaus of Astellas, Basiela, Gilead Sciences, Merck/MSD, and Pfizer, received research grants from Gilead Sciences and Pfizer and travel grants from Astellas, Merck/MSD, and Pfizer, and is a consultant to Basiela, Gilead Sciences, and MSD. U.G. received grants from Gilead Sciences and travel grants and speakers' fees from Pfizer and Gilead. O.B. received research grants from Pfizer and Gilead Sciences.

## REFERENCES

- Kousha M, Tadi R, Soubani AO. 2011. Pulmonary aspergillosis: a clinical review. *Eur Respir Rev* 20:156–174. <https://doi.org/10.1183/09059180.00001011>.
- van de Peppel RJ, von dem Borne PA, le Cessie S, de Boer MGJ. 2017. A new time-dependent approach for assessment of the impact of invasive aspergillosis shows effect on short- but not on long-term survival of patients with AML or high-risk MDS. *Bone Marrow Transplant* 52:883–888. <https://doi.org/10.1038/bmt.2017.71>.
- Snelders E, van der Lee HA, Kuijpers J, Rijs AJ, Varga J, Samson RA, Mellado E, Donders AR, Melchers WJ, Verweij PE. 2008. Emergence of azole resistance in *Aspergillus fumigatus* and spread of a single resistance mechanism. *PLoS Med* 5:e219. <https://doi.org/10.1371/journal.pmed.0050219>.
- Verweij PE, Chowdhary A, Melchers WJ, Meis JF. 2016. Azole resistance in *Aspergillus fumigatus*: can we retain the clinical use of mold-active antifungal azoles? *Clin Infect Dis* 62:362–368. <https://doi.org/10.1093/cid/civ885>.
- Wiederhold NP, Patterson TF. 2015. Emergence of azole resistance in *Aspergillus*. *Semin Respir Crit Care Med* 36:673–680. <https://doi.org/10.1055/s-0035-1562894>.
- Luo CX, Schnabel G. 2008. The cytochrome P450 lanosterol 14alpha-demethylase gene is a demethylation inhibitor fungicide resistance determinant in *Monilinia fructicola* field isolates from Georgia. *Appl Environ Microbiol* 74:359–366. <https://doi.org/10.1128/AEM.02159-07>.
- Ma Z, Proffer TJ, Jacobs JL, Sundin GW. 2006. Overexpression of the 14alpha-demethylase target gene (*cyp51*) mediates fungicide resistance in *Blumeriella jaapii*. *Appl Environ Microbiol* 72:2581–2585. <https://doi.org/10.1128/AEM.72.4.2581-2585.2006>.
- Parker JE, Warrilow AG, Price CL, Mullins JG, Kelly DE, Kelly SL. 2014. Resistance to antifungals that target CYP51. *J Chem Biol* 7:143–161. <https://doi.org/10.1007/s12154-014-0121-1>.
- Price CL, Parker JE, Warrilow AG, Kelly DE, Kelly SL. 2015. Azole fungicides—understanding resistance mechanisms in agricultural fungal pathogens. *Pest Manag Sci* 71:1054–1058. <https://doi.org/10.1002/ps.4029>.
- FRAC. 2017. Mechanisms of fungicide resistance. <http://www.frac.info/resistance-overview/mechanisms-of-fungicide-resistance>.
- Becher R, Wirsig SG. 2012. Fungal cytochrome P450 sterol 14alpha-demethylase (*CYP51*) and azole resistance in plant and human pathogens. *Appl Microbiol Biotechnol* 95:825–840. <https://doi.org/10.1007/s00253-012-4195-9>.
- Cools HJ, Fraaije BA. 2013. Update on mechanisms of azole resistance in *Mycosphaerella graminicola* and implications for future control. *Pest Manag Sci* 69:150–155. <https://doi.org/10.1002/ps.3348>.
- Verweij PE, Ananda-Rajah M, Andes D, Arendrup MC, Bruggemann RJ, Chowdhary A, Cornely OA, Denning DW, Groll AH, Izumikawa K, Kullberg BJ, Lagrou K, Maertens J, Meis JF, Newton P, Page I, Seyedmousavi S, Sheppard DC, Viscoli C, Warris A, Donnelly JP. 2015. International expert opinion on the management of infection caused by azole-resistant *Aspergillus fumigatus*. *Drug Resist Updat* 21–22:30–40. <https://doi.org/10.1016/j.drug.2015.08.001>.
- Verweij PE, Lestrade PP, Melchers WJ, Meis JF. 2016. Azole resistance surveillance in *Aspergillus fumigatus*: beneficial or biased? *J Antimicrob Chemother* 71:2079–2082. <https://doi.org/10.1093/jac/dkw259>.
- Patterson TF, Thompson GR, III, Denning DW, Fishman JA, Hadley S, Herbrecht R, Kontoyannis DP, Marr KA, Morrison VA, Nguyen MH, Segal BH, Steinbach WJ, Stevens DA, Walsh TJ, Wingard JR, Young JA, Bennett JE. 2016. Executive summary: practice guidelines for the diagnosis and management of aspergillosis: 2016 update by the Infectious Diseases Society of America. *Clin Infect Dis* 63:433–442. <https://doi.org/10.1093/cid/ciw444>.
- Bader O, Weig M, Reichard U, Lugert R, Kuhns M, Christner M, Held J, Peter S, Schumacher U, Buchheidt D, Tintelnot K, Gross U, MykoLabNet-D Partners. 2013. *cyp51A*-based mechanisms of *Aspergillus fumigatus* azole drug resistance present in clinical samples from Germany. *Antimicrob Agents Chemother* 57:3513–3517. <https://doi.org/10.1128/AAC.00167-13>.



17. Fraczek MG, Bromley M, Bueid A, Moore CB, Rajendran R, Rautemaa R, Ramage G, Denning DW, Bowyer P. 2013. The *cdr1B* efflux transporter is associated with non-*cyp51A*-mediated itraconazole resistance in *Aspergillus fumigatus*. *J Antimicrob Chemother* 68:1486–1496. <https://doi.org/10.1093/jac/dkt075>.
18. Bueid A, Howard SJ, Moore CB, Richardson MD, Harrison E, Bowyer P, Denning DW. 2010. Azole antifungal resistance in *Aspergillus fumigatus*: 2008 and 2009. *J Antimicrob Chemother* 65:2116–2118. <https://doi.org/10.1093/jac/dkq279>.
19. Camps SM, Dutilh BE, Arendrup MC, Rijs AJ, Snelders E, Huynen MA, Verweij PE, Melchers WJ. 2012. Discovery of a HapE mutation that causes azole resistance in *Aspergillus fumigatus* through whole genome sequencing and sexual crossing. *PLoS One* 7:e50034. <https://doi.org/10.1371/journal.pone.0050034>.
20. Gsaller F, Hortschansky P, Furukawa T, Carr PD, Rash B, Capilla J, Muller C, Bracher F, Bowyer P, Haas H, Brakhage AA, Bromley MJ. 2016. Sterol biosynthesis and azole tolerance is governed by the opposing actions of SrbA and the CCAAT binding complex. *PLoS Pathog* 12:e1005775. <https://doi.org/10.1371/journal.ppat.1005775>.
21. Paul S, Diekema D, Moyer-Rowley WS. 2013. Contributions of *Aspergillus fumigatus* ATP-binding cassette transporter proteins to drug resistance and virulence. *Eukaryot Cell* 12:1619–1628. <https://doi.org/10.1128/EC.00171-13>.
22. Meneau I, Coste AT, Sanglard D. 2016. Identification of *Aspergillus fumigatus* multidrug transporter genes and their potential involvement in antifungal resistance. *Med Mycol* 54:616–627. <https://doi.org/10.1093/mmy/myw005>.
23. Moyer-Rowley WS. 2015. Multiple mechanisms contribute to the development of clinically significant azole resistance in *Aspergillus fumigatus*. *Front Microbiol* 6:70. <https://doi.org/10.3389/fmicb.2015.00070>.
24. Steinmann J, Hamprecht A, Vehreschild MJ, Cornely OA, Buchheidt D, Spiess B, Koldehoff M, Buer J, Meis JF, Rath PM. 2015. Emergence of azole-resistant invasive aspergillosis in HSCT recipients in Germany. *J Antimicrob Chemother* 70:1522–1526. <https://doi.org/10.1093/jac/dku566>.
25. Chong GM, van der Beek MT, von dem Borne PA, Boelens J, Steel E, Kampinga GA, Span LF, Lagrou K, Maertens JA, Dingemans GJ, Gaajetaan GR, van Tegelen DW, Cornelissen JJ, Vonk AG, Rijnders BJ. 2016. PCR-based detection of *Aspergillus fumigatus* Cyp51A mutations on bronchoalveolar lavage: a multicentre validation of the AsperGenius assay(R) in 201 patients with haematological disease suspected for invasive aspergillosis. *J Antimicrob Chemother* 71:3528–3535. <https://doi.org/10.1093/jac/dkw323>.
26. van Paassen J, Russcher A, In't Veld-van Wingerden AW, Verweij PE, Kuijper EJ. 2016. Emerging aspergillosis by azole-resistant *Aspergillus fumigatus* at an intensive care unit in the Netherlands, 2010 to 2013. *Euro Surveill* 21:30300. <https://doi.org/10.2807/1560-7917.ES.2016.21.30.30300>.
27. Rössler S, Bader O, Stölzel F, Sommer U, Spiess B, Geibel S, Buchheidt D, Gross U, Baretton G, Jacobs E. 2017. Progressive dispersion of azole resistance in *Aspergillus fumigatus*: fatal invasive aspergillosis in a patient with acute myeloid leukemia infected with an A. *fumigatus* strain with a *cyp51A* TR<sub>46</sub> Y121F M172I T289A allele. *Antimicrob Agents Chemother* 61:e00270-17. <https://doi.org/10.1128/AAC.00270-17>.
28. Alanio A, Denis B, Hamane S, Raffoux E, Peffault de la Tour R, Touratier S, Bergeron A, Bretagne S. 2016. New therapeutic strategies for invasive aspergillosis in the era of azole resistance: how should the prevalence of azole resistance be defined? *J Antimicrob Chemother* 71:2075–2078. <https://doi.org/10.1093/jac/dkw036>.
29. Mair W, Lopez-Ruiz F, Stammler G, Clark W, Burnett F, Hollomon D, Ishii H, Thind TS, Brown JK, Fraaije B, Cools H, Shaw M, Fillinger S, Walker AS, Mellado E, Schnabel G, Mehl A, Oliver RP. 2016. Proposal for a unified nomenclature for target-site mutations associated with resistance to fungicides. *Pest Manag Sci* 72:1449–1459. <https://doi.org/10.1002/ps.4301>.
30. Walsh TJ, Anaissie EJ, Denning DW, Herbrecht R, Kontoyiannis DP, Marr KA, Morrison VA, Segal BH, Steinbach WJ, Stevens DA, van Burik JA, Wingard JR, Patterson TF, Infectious Diseases Society of America. 2008. Treatment of aspergillosis: clinical practice guidelines of the Infectious Diseases Society of America. *Clin Infect Dis* 46:327–360. <https://doi.org/10.1086/525258>.
31. Patterson TF, Thompson GR, III, Denning DW, Fishman JA, Hadley S, Herbrecht R, Kontoyiannis DP, Marr KA, Morrison VA, Nguyen MH, Segal BH, Steinbach WJ, Stevens DA, Walsh TJ, Wingard JR, Young JA, Bennett JE. 2016. Practice guidelines for the diagnosis and management of aspergillosis: 2016 update by the Infectious Diseases Society of America. *Clin Infect Dis* 63:e1–e60. <https://doi.org/10.1093/cid/ciw326>.
32. Munoz P, Alcalá L, Sánchez Conde M, Palomo J, Yanez J, Peláez T, Bouza E. 2003. The isolation of *Aspergillus fumigatus* from respiratory tract specimens in heart transplant recipients is highly predictive of invasive aspergillosis. *Transplantation* 75:326–329. <https://doi.org/10.1097/01.TP.0000044358.99414.B8>.
33. Horvath JA, Dummer S. 1996. The use of respiratory-tract cultures in the diagnosis of invasive pulmonary aspergillosis. *Am J Med* 100:171–178. [https://doi.org/10.1016/S0002-9343\(97\)89455-7](https://doi.org/10.1016/S0002-9343(97)89455-7).
34. Kwon-Chung KJ, Sugui JA. 2013. *Aspergillus fumigatus*—what makes the species a ubiquitous human fungal pathogen? *PLoS Pathog* 9:e1003743. <https://doi.org/10.1371/journal.ppat.1003743>.
35. Chowdhary A, Kathuria S, Xu J, Sharma C, Sundar G, Singh PK, Gaur SN, Hagen F, Klaassen CH, Meis JF. 2012. Clonal expansion and emergence of environmental multiple-triazole-resistant *Aspergillus fumigatus* strains carrying the TR(34)/L98H mutations in the *cyp51A* gene in India. *PLoS One* 7:e52871. <https://doi.org/10.1371/journal.pone.0052871>.
36. Dunne K, Hagen F, Pomeroy N, Meis JF, Rogers TR. 29 March 2017. Inter-country transfer of triazole-resistant *Aspergillus fumigatus* on plant bulbs. *Clin Infect Dis* <https://doi.org/10.1093/cid/cix257>.
37. Ahmad S, Joseph L, Hagen F, Meis JF, Khan Z. 2015. Concomitant occurrence of itraconazole-resistant and -susceptible strains of *Aspergillus fumigatus* in routine cultures. *J Antimicrob Chemother* 70: 412–415. <https://doi.org/10.1093/jac/dku410>.
38. Kolwijck E, van der Hoeven H, de Sevaux RG, ten Oever J, Rijstenberg LL, van der Lee HA, Zoll J, Melchers WJ, Verweij PE. 2016. Voriconazole-susceptible and voriconazole-resistant *Aspergillus fumigatus* coinfection. *Am J Respir Crit Care Med* 193:927–929. <https://doi.org/10.1164/rccm.201510-2104LE>.
39. Bader O, Tunnermann J, Dudakova A, Tangwattanachuleeporn M, Weig M, Gross U. 2015. Environmental isolates of azole-resistant *Aspergillus fumigatus* in Germany. *Antimicrob Agents Chemother* 59:4356–4359. <https://doi.org/10.1128/AAC.00100-15>.
40. Tangwattanachuleeporn M, Minarin N, Saichan S, Semsri P, Mitkornburee R, Gross U, Chindamporn A, Bader O. 2017. Prevalence of azole-resistant *Aspergillus fumigatus* in the environment of Thailand. *Med Mycol* 55:429–435. <https://doi.org/10.1093/mmy/myw090>.
41. Le Pape P, Laverne RA, Morio F, Alvarez-Moreno C. 2016. Multiple fungicide-driven alterations in azole-resistant *Aspergillus fumigatus*, Colombia, 2015. *Emerg Infect Dis* 22:156–157. <https://doi.org/10.3201/eid2201.150978>.
42. Prigitano A, Venier V, Cogliati M, De Lorenzis G, Esposto MC, Tortorano AM. 2014. Azole-resistant *Aspergillus fumigatus* in the environment of northern Italy, May 2011 to June 2012. *Euro Surveill* 19:20747. <https://doi.org/10.2807/1560-7917.ES.2014.19.12.20747>.
43. Chowdhary A, Sharma C, van den Boom M, Yntema JB, Hagen F, Verweij PE, Meis JF. 2014. Multi-azole-resistant *Aspergillus fumigatus* in the environment in Tanzania. *J Antimicrob Chemother* 69:2979–2983. <https://doi.org/10.1093/jac/dku259>.
44. Rocchi S, Daguindau E, Grenouillet F, Deconinck E, Bellanger AP, Garcia-Hermoso D, Bretagne S, Reboux G, Millon L. 2014. Azole-resistant *Aspergillus fumigatus* isolate with the TR<sub>34</sub>/L98H mutation in both a fungicide-sprayed field and the lung of a hematopoietic stem cell transplant recipient with invasive aspergillosis. *J Clin Microbiol* 52: 1724–1726. <https://doi.org/10.1128/JCM.03182-13>.
45. van der Linden JW, Camps SM, Kampinga GA, Arends JP, Debets-Ossenkopp YJ, Haas PJ, Rijnders BJ, Kuijper EJ, van Tiel FH, Varga J, Karawajczyk A, Zoll J, Melchers WJ, Verweij PE. 2013. Aspergillosis due to voriconazole highly resistant *Aspergillus fumigatus* and recovery of genetically related resistant isolates from domiciles. *Clin Infect Dis* 57:513–520. <https://doi.org/10.1093/cid/cit320>.
46. Ahmad S, Khan Z, Hagen F, Meis JF. 2014. Occurrence of triazole-resistant *Aspergillus fumigatus* with TR<sub>34</sub>/L98H mutations in outdoor and hospital environment in Kuwait. *Environ Res* 133:20–26. <https://doi.org/10.1016/j.envres.2014.05.009>.
47. Bhabhra R, Askew DS. 2005. Thermotolerance and virulence of *Aspergillus fumigatus*: role of the fungal nucleolus. *Med Mycol* 43(Suppl 1):S87–S93. <https://doi.org/10.1080/13693780400029486>.
48. van der Linden JWM, Arendrup MC, van der Lee HAL, Melchers WJG, Verweij PE. 2009. Azole containing agar plates as a screening tool for azole resistance of *Aspergillus fumigatus*. *Mycoses* 52:19–28. <https://doi.org/10.1111/j.1439-0507.2009.01781.x>.

49. Mortensen KL, Mellado E, Lass-Flörl C, Rodriguez-Tudela JL, Johansen HK, Arendrup MC. 2010. Environmental study of azole-resistant *Aspergillus fumigatus* and other aspergilli in Austria, Denmark, and Spain. *Antimicrob Agents Chemother* 54:4545–4549. <https://doi.org/10.1128/AAC.00692-10>.
50. Snelders E, Huis In't Veld RA, Rijs AJ, Kema GH, Melchers WJ, Verweij PE. 2009. Possible environmental origin of resistance of *Aspergillus fumigatus* to medical triazoles. *Appl Environ Microbiol* 75:4053–4057. <https://doi.org/10.1128/AEM.00231-09>.
51. Edlind TD, Henry KW, Metera KA, Katiyar SK. 2001. *Aspergillus fumigatus* cyp51 sequence: potential basis for fluconazole resistance. *Med Mycol* 39:299–302.
52. Leonardelli F, Macedo D, Dudiuk C, Cabeza MS, Gamarra S, Garcia-Effron G. 2016. *Aspergillus fumigatus* intrinsic fluconazole resistance is due to the naturally occurring T301I substitution in Cyp51A. *Antimicrob Agents Chemother* 60:5420–5426. <https://doi.org/10.1128/AAC.00905-16>.
53. Warrilow AG, Melo N, Martel CM, Parker JE, Nes WD, Kelly SL, Kelly DE. 2010. Expression, purification, and characterization of *Aspergillus fumigatus* sterol 14-alpha demethylase (CYP51) isoenzymes A and B. *Antimicrob Agents Chemother* 54:4225–4234. <https://doi.org/10.1128/AAC.00316-10>.
54. Mortensen KL, Jensen RH, Johansen HK, Skov M, Pressler T, Howard SJ, Leatherbarrow H, Mellado E, Arendrup MC. 2011. *Aspergillus* species and other molds in respiratory samples from patients with cystic fibrosis: a laboratory-based study with focus on *Aspergillus fumigatus* azole resistance. *J Clin Microbiol* 49:2243–2251. <https://doi.org/10.1128/JCM.00213-11>.
55. Dannaoui E, Meletiadis J, Tortorano AM, Symoens F, Nolard N, Viviani MA, Piens MA, Lebeau B, Verweij PE, Grillot R, EBGA Network. 2004. Susceptibility testing of sequential isolates of *Aspergillus fumigatus* recovered from treated patients. *J Med Microbiol* 53:129–134. <https://doi.org/10.1099/jmm.0.05326-0>.
56. Howard SJ, Cerar D, Anderson MJ, Albarrag A, Fisher MC, Pasqualotto AC, Laverdiere M, Arendrup MC, Perlin DS, Denning DW. 2009. Frequency and evolution of azole resistance in *Aspergillus fumigatus* associated with treatment failure. *Emerg Infect Dis* 15:1068–1076. <https://doi.org/10.3201/eid1507.090043>.
57. Rodriguez-Tudela JL, Alcazar-Fuoli L, Mellado E, Alastruey-Izquierdo A, Monzon A, Cuenca-Estrella M. 2008. Epidemiological cutoffs and cross-resistance to azole drugs in *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 52:2468–2472. <https://doi.org/10.1128/AAC.00156-08>.
58. Hamprecht A, Buchheidt D, Vehreschild JJ, Cornely OA, Spiess B, Plum G, Halbsguth TV, Kutsch N, Stippel D, Kahl P, Persigehl T, Steinbach A, Bos B, Hallek M, Vehreschild MJ. 2012. Azole-resistant invasive aspergillosis in a patient with acute myeloid leukaemia in Germany. *Euro Surveill* 17:20262.
59. Pelaez T, Monteiro MC, Garcia-Rubio R, Bouza E, Gomez-Lopez A, Mellado E. 2015. First detection of *Aspergillus fumigatus* azole-resistant strain due to Cyp51A TR<sub>46</sub>/Y121F/T289A in an azole-naïve patient in Spain. *New Microbes New Infect* 6:33–34. <https://doi.org/10.1016/j.nmni.2015.04.005>.
60. Camps SM, van der Linden JW, Li Y, Kuijper EJ, van Dissel JT, Verweij PE, Melchers WJ. 2012. Rapid induction of multiple resistance mechanisms in *Aspergillus fumigatus* during azole therapy: a case study and review of the literature. *Antimicrob Agents Chemother* 56:10–16. <https://doi.org/10.1128/AAC.05088-11>.
61. Morio F, Aubin GG, Danner-Boucher I, Haloun A, Sacchetto E, Garcia-Hermoso D, Bretagne S, Miegerville M, Le Pape P. 2012. High prevalence of triazole resistance in *Aspergillus fumigatus*, especially mediated by TR/L98H, in a French cohort of patients with cystic fibrosis. *J Antimicrob Chemother* 67:1870–1873. <https://doi.org/10.1093/jac/dks160>.
62. Alanio A, Cabaret O, Sitterle E, Costa JM, Brisse S, Cordonnier C, Bretagne S. 2012. Azole preexposure affects the *Aspergillus fumigatus* population in patients. *Antimicrob Agents Chemother* 56:4948–4950. <https://doi.org/10.1128/AAC.05990-11>.
63. Fischer J, van Koningsbruggen-Rietschel S, Rietschel E, Vehreschild MJ, Wisplinghoff H, Kronke M, Hamprecht A. 2014. Prevalence and molecular characterization of azole resistance in *Aspergillus* spp. isolates from German cystic fibrosis patients. *J Antimicrob Chemother* 69:1533–1536. <https://doi.org/10.1093/jac/dku009>.
64. Lavergne RA, Morio F, Favennec L, Dominique S, Meis JF, Gargala G, Verweij PE, Le Pape P. 2015. First description of azole-resistant *Aspergillus fumigatus* due to TR<sub>46</sub>/Y121F/T289A mutation in France. *Antimicrob Agents Chemother* 59:4331–4335. <https://doi.org/10.1128/AAC.00127-15>.
65. Lazzarini C, Esposto MC, Prigitano A, Cogliati M, De Lorenzis G, Tortorano AM. 2015. Azole resistance in *Aspergillus fumigatus* clinical isolates from an Italian culture collection. *Antimicrob Agents Chemother* 60:682–685. <https://doi.org/10.1128/AAC.02234-15>.
66. van Ingen J, van der Lee HA, Rijs TA, Zoll J, Leenstra T, Melchers WJ, Verweij PE. 2015. Azole, polyene and echinocandin MIC distributions for wild-type, TR<sub>34</sub>/L98H and TR<sub>46</sub>/Y121F/T289A *Aspergillus fumigatus* isolates in the Netherlands. *J Antimicrob Chemother* 70:178–181. <https://doi.org/10.1093/jac/dku364>.
67. van Ingen J, van der Lee HA, Rijs AJ, Snelders E, Melchers WJ, Verweij PE. 2015. High-level pan-azole-resistant aspergillosis. *J Clin Microbiol* 53:2343–2345. <https://doi.org/10.1128/JCM.00502-15>.
68. Astvad KM, Jensen RH, Hassan TM, Mathiasen EG, Thomsen GM, Pedersen UG, Christensen M, Hilberg O, Arendrup MC. 2014. First detection of TR<sub>46</sub>/Y121F/T289A and TR<sub>34</sub>/L98H alterations in *Aspergillus fumigatus* isolates from azole-naïve patients in Denmark despite negative findings in the environment. *Antimicrob Agents Chemother* 58:5096–5101. <https://doi.org/10.1128/AAC.02855-14>.
69. Choukri F, Botterel F, Sitterle E, Bassinet L, Foulet F, Guillot J, Costa JM, Fauchet N, Dannaoui E. 2015. Prospective evaluation of azole resistance in *Aspergillus fumigatus* clinical isolates in France. *Med Mycol* 53: 593–596. <https://doi.org/10.1093/mmy/myv029>.
70. Liu M, Zeng R, Zhang L, Li D, Lv G, Shen Y, Zheng H, Zhang Q, Zhao J, Zheng N, Liu W. 2015. Multiple cyp51A-based mechanisms identified in azole-resistant isolates of *Aspergillus fumigatus* from China. *Antimicrob Agents Chemother* 59:4321–4325. <https://doi.org/10.1128/AAC.00003-15>.
71. Mohammadi F, Hashemi SJ, Zoll J, Melchers WJ, Rafati H, Dehghan P, Rezaie S, Tolooe A, Tamadon Y, van der Lee HA, Verweij PE, Seyedmousavi S. 2015. Quantitative analysis of single-nucleotide polymorphism for rapid detection of TR<sub>34</sub>/L98H- and TR<sub>46</sub>/Y121F/T289A-positive *Aspergillus fumigatus* isolates obtained from patients in Iran from 2010 to 2014. *Antimicrob Agents Chemother* 60:387–392. <https://doi.org/10.1128/AAC.02326-15>.
72. Jensen RH, Hagen F, Astvad KM, Tyrone A, Meis JF, Arendrup MC. 2016. Azole-resistant *Aspergillus fumigatus* in Denmark: a laboratory-based study on resistance mechanisms and genotypes. *Clin Microbiol Infect* 22:570.e1–570.e9. <https://doi.org/10.1016/j.cmi.2016.04.001>.
73. Chen Y, Lu Z, Zhao J, Zou Z, Gong Y, Qu F, Bao Z, Qiu G, Song M, Zhang Q, Liu L, Hu M, Han X, Tian S, Zhao J, Chen F, Zhang C, Sun Y, Verweij PE, Huang L, Han L. 2016. Epidemiology and molecular characterizations of azole resistance in clinical and environmental *Aspergillus fumigatus* isolates from China. *Antimicrob Agents Chemother* 60: 5878–5884. <https://doi.org/10.1128/AAC.01005-16>.
74. Howard SJ, Pasqualotto AC, Denning DW. 2010. Azole resistance in allergic bronchopulmonary aspergillosis and *Aspergillus* bronchitis. *Clin Microbiol Infect* 16:683–688. <https://doi.org/10.1111/j.1469-0691.2009.02911.x>.
75. Albarrag AM, Anderson MJ, Howard SJ, Robson GD, Warn PA, Sanglard D, Denning DW. 2011. Interrogation of related clinical pan-azole-resistant *Aspergillus fumigatus* strains: G138C, Y431C, and G434C single nucleotide polymorphisms in cyp51A, upregulation of cyp51A, and integration and activation of transposon Atf1 in the cyp51A promoter. *Antimicrob Agents Chemother* 55:5113–5121. <https://doi.org/10.1128/AAC.00517-11>.
76. Mushi MF, Buname G, Bader O, Gross U, Mshana SE. 2016. *Aspergillus fumigatus* carrying TR<sub>34</sub>/L98H resistance allele causing complicated suppurative otitis media in Tanzania: call for improved diagnosis of fungi in sub-Saharan Africa. *BMC Infect Dis* 16:464. <https://doi.org/10.1186/s12879-016-1796-4>.
77. Snelders E, Karawajczyk A, Verhoeven RJ, Venselaar H, Schaftenaar G, Verweij PE, Melchers WJ. 2011. The structure-function relationship of the *Aspergillus fumigatus* Cyp51A L98H conversion by site-directed mutagenesis: the mechanism of L98H azole resistance. *Fungal Genet Biol* 48:1062–1070. <https://doi.org/10.1016/j.fgb.2011.08.002>.
78. Jeanvoine A, Rocchi S, Reboux G, Crini N, Crini G, Millon L. 12 May 2017. Azole-resistant *Aspergillus fumigatus* in sawmills of Eastern France. *J Appl Microbiol* <https://doi.org/10.1111/jam.13488>.
79. Nawrot U, Kurzyk E, Arendrup MC, Mroczynska M, Włodarczyk K, Sulik-Tyska B, Wroblewska M, Ussowicz M, Zdziarski P, Niewinska K, Brillowska-Dabrowska A. 4 March 2017. Detection of Polish clinical

- Aspergillus fumigatus* isolates resistant to triazoles. *Med Mycol* <https://doi.org/10.1093/mmy/myx012>.
80. Zhang M, Feng CL, Chen F, He Q, Su X, Shi Y. 2017. Triazole resistance in *Aspergillus fumigatus* clinical isolates obtained in Nanjing, China. *Chin Med J (Engl)* 130:665–668. <https://doi.org/10.4103/0366-6999.201609>.
  81. Zhang J, Snelders E, Zwaan BJ, Schoustra SE, Meis JF, van Dijk K, Hagen F, van der Beek MT, Kampinga GA, Zoll J, Melchers WJG, Verweij PE, Debets AJM. 2017. A novel environmental azole resistance mutation in *Aspergillus fumigatus* and a possible role of sexual reproduction in its emergence. *mBio* 8:e00791-17. <https://doi.org/10.1128/mBio.00791-17>.
  82. Snelders E, Karawajczyk A, Schaftenaar G, Verweij PE, Melchers WJ. 2010. Azole resistance profile of amino acid changes in *Aspergillus fumigatus* CYP51A based on protein homology modeling. *Antimicrob Agents Chemother* 54:2425–2430. <https://doi.org/10.1128/AAC.01599-09>.
  83. Garcia-Effron G, Dilger A, Alcazar-Fuoli L, Park S, Mellado E, Perlin DS. 2008. Rapid detection of triazole antifungal resistance in *Aspergillus fumigatus*. *J Clin Microbiol* 46:1200–1206. <https://doi.org/10.1128/JCM.02330-07>.
  84. Mellado E, Garcia-Effron G, Alcazar-Fuoli L, Melchers WJ, Verweij PE, Cuenca-Estrella M, Rodriguez-Tudela JL. 2007. A new *Aspergillus fumigatus* resistance mechanism conferring in vitro cross-resistance to azole antifungals involves a combination of *cyp51A* alterations. *Antimicrob Agents Chemother* 51:1897–1904. <https://doi.org/10.1128/AAC.01092-06>.
  85. Mavridou E, Bruggemann RJ, Melchers WJ, Mouton JW, Verweij PE. 2010. Efficacy of posaconazole against three clinical *Aspergillus fumigatus* isolates with mutations in the *cyp51A* gene. *Antimicrob Agents Chemother* 54:860–865. <https://doi.org/10.1128/AAC.00931-09>.
  86. van der Linden JW, Jansen RR, Bresters D, Visser CE, Geerlings SE, Kuijper EJ, Melchers WJ, Verweij PE. 2009. Azole-resistant central nervous system aspergillosis. *Clin Infect Dis* 48:1111–1113. <https://doi.org/10.1086/597465>.
  87. Lockhart SR, Frade JP, Etienne KA, Pfaller MA, Diekema DJ, Balajee SA. 2011. Azole resistance in *Aspergillus fumigatus* isolates from the ARTEMIS global surveillance study is primarily due to the TR/L98H mutation in the *cyp51A* gene. *Antimicrob Agents Chemother* 55:4465–4468. <https://doi.org/10.1128/AAC.00185-11>.
  88. Burgel PR, Baixench MT, Amsellem M, Audureau E, Chapron J, Kanaan R, Honore I, Dupouy-Camet J, Dusser D, Klaassen CH, Meis JF, Hubert D, Paugam A. 2012. High prevalence of azole-resistant *Aspergillus fumigatus* in adults with cystic fibrosis exposed to itraconazole. *Antimicrob Agents Chemother* 56:869–874. <https://doi.org/10.1128/AAC.05077-11>.
  89. Rath PM, Buchheidt D, Spiess B, Arfanis E, Buer J, Steinmann J. 2012. First reported case of azole-resistant *Aspergillus fumigatus* due to the TR/L98H mutation in Germany. *Antimicrob Agents Chemother* 56:6060–6061. <https://doi.org/10.1128/AAC.01017-12>.
  90. Tashiro M, Izumikawa K, Minematsu A, Hirano K, Iwanaga N, Ide S, Mihara T, Hosogaya N, Takazono T, Morinaga Y, Nakamura S, Kurihara S, Imamura Y, Miyazaki T, Nishino T, Tsukamoto M, Kakeya H, Yamamoto Y, Yanagihara K, Yasuoka A, Tashiro T, Kohno S. 2012. Antifungal susceptibilities of *Aspergillus fumigatus* clinical isolates obtained in Nagasaki, Japan. *Antimicrob Agents Chemother* 56:584–587. <https://doi.org/10.1128/AAC.05394-11>.
  91. van der Linden JW, Snelders E, Kampinga GA, Rijnders BJ, Mattsson E, Debets-Ossenkopp YJ, Kuijper EJ, Van Tiel FH, Melchers WJ, Verweij PE. 2011. Clinical implications of azole resistance in *Aspergillus fumigatus*, The Netherlands, 2007–2009. *Emerg Infect Dis* 17:1846–1854. <https://doi.org/10.3201/eid1710.110226>.
  92. Chowdhary A, Kathuria S, Randhawa HS, Gaur SN, Klaassen CH, Meis JF. 2012. Isolation of multiple-triazole-resistant *Aspergillus fumigatus* strains carrying the TR/L98H mutations in the *cyp51A* gene in India. *J Antimicrob Chemother* 67:362–366. <https://doi.org/10.1093/jac/dkr443>.
  93. Chowdhary A, Sharma C, Kathuria S, Hagen F, Meis JF. 2015. Prevalence and mechanism of triazole resistance in *Aspergillus fumigatus* in a referral chest hospital in Delhi, India and an update of the situation in Asia. *Front Microbiol* 6:428. <https://doi.org/10.3389/fmicb.2015.00428>.
  94. Gregson L, Goodwin J, Johnson A, McEntee L, Moore CB, Richardson M, Hope WW, Howard SJ. 2013. In vitro susceptibility of *Aspergillus fumigatus* to isavuconazole: correlation with itraconazole, voriconazole, and posaconazole. *Antimicrob Agents Chemother* 57:5778–5780. <https://doi.org/10.1128/AAC.01141-13>.
  95. Wiederhold NP, Gil VG, Gutierrez F, Lindner JR, Albataineh MT, McCarthy DI, Sanders C, Fan H, Fothergill AW, Sutton DA. 2016. First detection of TR<sub>34</sub> L98H and TR<sub>46</sub> Y121F/T289A *cyp51* mutations in *Aspergillus fumigatus* isolates in the United States. *J Clin Microbiol* 54:168–171. <https://doi.org/10.1128/JCM.02478-15>.
  96. Wu CJ, Wang HC, Lee JC, Lo HJ, Dai CT, Chou PH, Ko WC, Chen YC. 2015. Azole-resistant *Aspergillus fumigatus* isolates carrying TR<sub>34</sub>/L98H mutations in Taiwan. *Mycoses* 58:544–549. <https://doi.org/10.1111/myc.12354>.
  97. Badali H, Vaezi A, Haghani I, Yazdanparast SA, Hedayati MT, Mousavi B, Ansari S, Hagen F, Meis JF, Chowdhary A. 2013. Environmental study of azole-resistant *Aspergillus fumigatus* with TR<sub>34</sub>/L98H mutations in the *cyp51A* gene in Iran. *Mycoses* 56:659–663. <https://doi.org/10.1111/myc.12089>.
  98. Vermeulen E, Maertens J, Schoemans H, Lagrou K. 2012. Azole-resistant *Aspergillus fumigatus* due to TR<sub>46</sub>/Y121F/T289A mutation emerging in Belgium, July 2012. *Euro Surveill* 17:20326.
  99. da Silva Ferreira ME, Capellaro JL, dos Reis Marques E, Malavazi I, Perlin D, Park S, Anderson JB, Colombo AL, Arthington-Skaggs BA, Goldman MH, Goldman GH. 2004. *In vitro* evolution of itraconazole resistance in *Aspergillus fumigatus* involves multiple mechanisms of resistance. *Antimicrob Agents Chemother* 48:4405–4413. <https://doi.org/10.1128/AAC.48.11.4405-4413.2004>.
  100. Diaz-Guerra TM, Mellado E, Cuenca-Estrella M, Rodriguez-Tudela JL. 2003. A point mutation in the 14 $\alpha$ -sterol demethylase gene *cyp51A* contributes to itraconazole resistance in *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 47:1120–1124. <https://doi.org/10.1128/AAC.47.3.1120-1124.2003>.
  101. Garcia-Effron G, Mellado E, Gomez-Lopez A, Alcazar-Fuoli L, Cuenca-Estrella M, Rodriguez-Tudela JL. 2005. Differences in interactions between azole drugs related to modifications in the 14- $\alpha$  sterol demethylase gene (*cyp51A*) of *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 49:2119–2121. <https://doi.org/10.1128/AAC.49.5.2119-2121.2005>.
  102. Mellado E, Garcia-Effron G, Alcazar-Fuoli L, Cuenca-Estrella M, Rodriguez-Tudela JL. 2004. Substitutions at methionine 220 in the 14 $\alpha$ -sterol demethylase (*Cyp51A*) of *Aspergillus fumigatus* are responsible for resistance *in vitro* to azole antifungal drugs. *Antimicrob Agents Chemother* 48:2747–2750. <https://doi.org/10.1128/AAC.48.7.2747-2750.2004>.
  103. Mann PA, Parmegiani RM, Wei SQ, Mendrick CA, Li X, Loebenberg D, DiDomenico B, Hare RS, Walker SS, McNicholas PM. 2003. Mutations in *Aspergillus fumigatus* resulting in reduced susceptibility to posaconazole appear to be restricted to a single amino acid in the cytochrome P450 14 $\alpha$ -demethylase. *Antimicrob Agents Chemother* 47:577–581. <https://doi.org/10.1128/AAC.47.2.577-581.2003>.
  104. Howard SJ, Webster I, Moore CB, Gardiner RE, Park S, Perlin DS, Denning DW. 2006. Multi-azole resistance in *Aspergillus fumigatus*. *Int J Antimicrob Agents* 28:450–453. <https://doi.org/10.1016/j.ijantimicag.2006.08.017>.
  105. Hagiwara D, Takahashi H, Fujimoto M, Sugahara M, Misawa Y, Gonoi T, Itoyama S, Watanabe A, Kamei K. 2016. Multi-azole resistant *Aspergillus fumigatus* harboring TR<sub>46</sub>/Y121F/T289A isolated in Japan. *J Infect Chemother* 22:577–579. <https://doi.org/10.1016/j.jiac.2016.01.015>.
  106. Leonardelli F, Theill L, Nardin ME, Macedo D, Dudiuk C, Mendez E, Gamarra S, Garcia-Effron G. 2017. First itraconazole resistant *Aspergillus fumigatus* clinical isolate harbouring a G54E substitution in *Cyp51A* in South America. *Rev Iberoam Micol* 34:46–48. <https://doi.org/10.1016/j.riam.2016.05.005>.
  107. Montesinos I, Argudin MA, Hites M, Ahajjam F, Dodemont M, Dagyan C, Bakali M, Etienne I, Jacobs F, Knoop C, Patteet S, Lagrou K. 2017. Culture-based methods and molecular tools for azole-resistant *Aspergillus fumigatus* detection in a Belgian university hospital. *J Clin Microbiol* 55:2391–2399. <https://doi.org/10.1128/JCM.00520-17>.
  108. Sharma C, Hagen F, Moroti R, Meis JF, Chowdhary A. 2015. Triazole-resistant *Aspergillus fumigatus* harbouring G54 mutation: is it de novo or environmentally acquired? *J Glob Antimicrob Resist* 3:69–74. <https://doi.org/10.1016/j.jgar.2015.01.005>.
  109. Rybak JM, Marx KR, Nishimoto AT, Rogers PD. 2015. Isavuconazole: pharmacology, pharmacodynamics, and current clinical experience with a new triazole antifungal agent. *Pharmacotherapy* 35:1037–1051. <https://doi.org/10.1002/phar.1652>.
  110. Balajee SA, Nickle D, Varga J, Marr KA. 2006. Molecular studies reveal



- frequent misidentification of *Aspergillus fumigatus* by morphotyping. Eukaryot Cell 5:1705–1712. <https://doi.org/10.1128/EC.00162-06>.
111. Balajee SA, Houbraeken J, Verweij PE, Hong SB, Yaghuchi T, Varga J, Samson RA. 2007. *Aspergillus* species identification in the clinical setting. Stud Mycol 59:39–46. <https://doi.org/10.3114/sim.2007.59.05>.
  112. De Hoog G, Guarro J, Figueras M. 2000. Atlas of clinical fungi, 2nd ed. American Society for Microbiology, Washington, DC.
  113. Faria-Ramos I, Farinha S, Neves-Maia J, Tavares PR, Miranda IM, Estevinho LM, Pina-Vaz C, Rodrigues AG. 2014. Development of cross-resistance by *Aspergillus fumigatus* to clinical azoles following exposure to prochloraz, an agricultural azole. BMC Microbiol 14:155. <https://doi.org/10.1186/1471-2180-14-155>.
  114. Samson RA, Varga J. 2009. What is a species in *Aspergillus*? Med Mycol 47(Suppl 1):S13–S20. <https://doi.org/10.1080/13693780802354011>.
  115. Kocsube S, Varga J. 2017. Targeting conserved genes in *Aspergillus* species. Methods Mol Biol 1542:131–140. [https://doi.org/10.1007/978-1-4939-6707-0\\_7](https://doi.org/10.1007/978-1-4939-6707-0_7).
  116. De Carolis E, Posteraro B, Lass-Flörl C, Vella A, Florio AR, Torelli R, Girmenia C, Colozza C, Tortorano AM, Sanguinetti M, Fadda G. 2012. Species identification of *Aspergillus*, *Fusarium* and *Mucorales* with direct surface analysis by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. Clin Microbiol Infect 18:475–484. <https://doi.org/10.1111/j.1469-0691.2011.03599.x>.
  117. Iriart X, Lavergne RA, Fillaux J, Valentin A, Magnaval JF, Berry A, Cassaing S. 2012. Routine identification of medical fungi by the new Vitek MS matrix-assisted laser desorption ionization-time of flight system with a new time-effective strategy. J Clin Microbiol 50:2107–2110. <https://doi.org/10.1128/JCM.06713-11>.
  118. Bille E, Dauphin B, Leto J, Bougnoux ME, Beretti JL, Lotz A, Suarez S, Meyer J, Join-Lambert O, Descamps P, Grall N, Mory F, Dubreuil L, Berche P, Nassif X, Ferroni A. 2012. MALDI-TOF MS Andromas strategy for the routine identification of bacteria, mycobacteria, yeasts, *Aspergillus* spp. and positive blood cultures. Clin Microbiol Infect 18: 1117–1125. <https://doi.org/10.1111/j.1469-0691.2011.03688.x>.
  119. Houbraeken J, Weig M, Gross U, Meijer M, Bader O. 2016. *Aspergillus oerlinghausenensis*, a new mould species closely related to *A. fumigatus*. FEMS Microbiol Lett 363:fnv236. <https://doi.org/10.1093/femsle/fnv236>.
  120. Lass-Flörl C. 2010. *In vitro* susceptibility testing in *Aspergillus* species: an update. Future Microbiol 5:789–799. <https://doi.org/10.2217/fmb.10.34>.
  121. Rex JH, Alexander BD, Andes D, Arthington-Skaggs B, Brown SD, Chaturvedi V, Espinel-Ingroff A, Ghanoum MA, Knapp CC, Motyl MR, Ostrosky-Zeichner L, Pfaller JB, Sheehan DJ, Walsh T. 2008. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi; approved standard—2nd ed. CLSI document M38-A2, 2nd ed, vol 28. CLSI, Wayne, PA.
  122. Espinel-Ingroff A, Fothergill A, Ghanoum M, Manavathu E, Ostrosky-Zeichner L, Pfaller M, Rinaldi M, Schell W, Walsh T. 2005. Quality control and reference guidelines for CLSI broth microdilution susceptibility method (M38-A document) for amphotericin B, itraconazole, posaconazole, and voriconazole. J Clin Microbiol 43:5243–5246. <https://doi.org/10.1128/JCM.43.10.5243-5246.2005>.
  123. Subcommittee on Antifungal Susceptibility Testing of the ESCMID European Committee for Antimicrobial Susceptibility Testing. 2008. EUCAST technical note on the method for the determination of broth dilution minimum inhibitory concentrations of antifungal agents for conidia-forming moulds. Clin Microbiol Infect 14:982–984. <https://doi.org/10.1111/j.1469-0691.2008.02086.x>.
  124. Espinel-Ingroff A. 2006. Comparison of three commercial assays and a modified disk diffusion assay with two broth microdilution reference assays for testing zygomycetes, *Aspergillus* spp., *Candida* spp., and *Cryptococcus neoformans* with posaconazole and amphotericin B. J Clin Microbiol 44:3616–3622. <https://doi.org/10.1128/JCM.01187-06>.
  125. Espinel-Ingroff A, Arthington-Skaggs B, Iqbal N, Ellis D, Pfaller MA, Messer S, Rinaldi M, Fothergill A, Gibbs DL, Wang A. 2007. Multicenter evaluation of a new disk agar diffusion method for susceptibility testing of filamentous fungi with voriconazole, posaconazole, itraconazole, amphotericin B, and caspofungin. J Clin Microbiol 45:1811–1820. <https://doi.org/10.1128/JCM.00134-07>.
  126. Szekely A, Johnson EM, Warnock DW. 1999. Comparison of E-test and broth microdilution methods for antifungal drug susceptibility testing of molds. J Clin Microbiol 37:1480–1483.
  127. Pfaller MA, Messer SA, Mills K, Bolmstrom A. 2000. *In vitro* susceptibility testing of filamentous fungi: comparison of Etest and reference microdilution methods for determining itraconazole MICs. J Clin Microbiol 38:3359–3361.
  128. Arendrup MC, Verweij P, Nielsen HV. 2017. Evaluation of MIC strip isavuconazole test for susceptibility testing of wild-type and non-wild-type *Aspergillus fumigatus* isolates. Antimicrob Agents Chemother 61:e01659-16. <https://doi.org/10.1128/AAC.01659-16>.
  129. Castro C, Serrano MC, Flores B, Espinel-Ingroff A, Martin-Mazuelos E. 2004. Comparison of the Sensititre YeastOne colorimetric antifungal panel with a modified NCCLS M38-A method to determine the activity of voriconazole against clinical isolates of *Aspergillus* spp. J Clin Microbiol 42:4358–4360. <https://doi.org/10.1128/JCM.42.9.4358-4360.2004>.
  130. Arendrup MC, Cuenca-Estrella M, Lass-Flörl C, Hope WW. 2013. Breakpoints for antifungal agents: an update from EUCAST focussing on echinocandins against *Candida* spp. and triazoles against *Aspergillus* spp. Drug Resist Updat 16:81–95. <https://doi.org/10.1016/j.drup.2014.01.001>.
  131. Lass-Flörl C, Perkhof S, Mayr A. 2010. *In vitro* susceptibility testing in fungi: a global perspective on a variety of methods. Mycoses 53:1–11. <https://doi.org/10.1111/j.1439-0507.2009.01813.x>.
  132. Mosquera J, Denning DW. 2002. Azole cross-resistance in *Aspergillus fumigatus*. Antimicrob Agents Chemother 46:556–557. <https://doi.org/10.1128/AAC.46.2.556-557.2002>.
  133. EUCAST. 2015. Antifungal agents. Breakpoint tables for interpretation of MICs. [http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST\\_files/AFST/Clinical\\_breakpoints/Antifungal\\_breakpoints\\_v\\_8.0\\_November\\_2015.pdf](http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/AFST/Clinical_breakpoints/Antifungal_breakpoints_v_8.0_November_2015.pdf). Accessed 5 January 2016.
  134. Rex JH, Pfaller MA. 2002. Has antifungal susceptibility testing come of age? Clin Infect Dis 35:982–989. <https://doi.org/10.1086/342384>.
  135. Vanden Bossche H, Koymans L, Moereels H. 1995. P450 inhibitors of use in medical treatment: focus on mechanisms of action. Pharmacol Ther 67:79–100. [https://doi.org/10.1016/0163-7258\(95\)00011-5](https://doi.org/10.1016/0163-7258(95)00011-5).
  136. Osherov N, Kontoyannis DP, Romans A, May GS. 2001. Resistance to itraconazole in *Aspergillus nidulans* and *Aspergillus fumigatus* is conferred by extra copies of the *A. nidulans* P-450 14 $\alpha$ -demethylase gene, *pdmA*. J Antimicrob Chemother 48:75–81. <https://doi.org/10.1093/jac/48.1.75>.
  137. Taylor JW. 2014. Evolutionary perspectives on human fungal pathogens. Cold Spring Harb Perspect Med 5:a019588. <https://doi.org/10.1101/cshperspect.a019588>.
  138. Hawkins NJ, Cools HJ, Sierotzki H, Shaw MW, Knogge W, Kelly SL, Kelly DE, Fraaije BA. 2014. Paralog re-emergence: a novel, historically contingent mechanism in the evolution of antimicrobial resistance. Mol Biol Evol 31:1793–1802. <https://doi.org/10.1093/molbev/msu134>.
  139. Mellado E, Diaz-Guerra TM, Cuenca-Estrella M, Rodriguez-Tudela JL. 2001. Identification of two different 14- $\alpha$  sterol demethylase-related genes (*cyp51A* and *cyp51B*) in *Aspergillus fumigatus* and other *Aspergillus* species. J Clin Microbiol 39:2431–2438. <https://doi.org/10.1128/JCM.39.7.2431-2438.2001>.
  140. Martel CM, Parker JE, Warrilow AG, Rolley NJ, Kelly SL, Kelly DE. 2010. Complementation of a *Saccharomyces cerevisiae* *ERG11/CYP51* (sterol 14 $\alpha$ -demethylase) doxycycline-regulated mutant and screening of the azole sensitivity of *Aspergillus fumigatus* isoenzymes CYP51A and CYP51B. Antimicrob Agents Chemother 54:4920–4923. <https://doi.org/10.1128/AAC.00349-10>.
  141. Warrilow AG, Parker JE, Price CL, Nes WD, Kelly SL, Kelly DE. 2015. *In vitro* biochemical study of CYP51-mediated azole resistance in *Aspergillus fumigatus*. Antimicrob Agents Chemother 59:7771–7778. <https://doi.org/10.1128/AAC.01806-15>.
  142. Flowers SA, Colon B, Whaley SG, Schuler MA, Rogers PD. 2015. Contribution of clinically derived mutations in *ERG11* to azole resistance in *Candida albicans*. Antimicrob Agents Chemother 59:450–460. <https://doi.org/10.1128/AAC.03470-14>.
  143. Xiang MJ, Liu JY, Ni PH, Wang S, Shi C, Wei B, Ni YX, Ge HL. 2013. Erg11 mutations associated with azole resistance in clinical isolates of *Candida albicans*. FEMS Yeast Res 13:386–393. <https://doi.org/10.1111/1567-1364.12042>.
  144. Hagiwara D, Watanabe A, Kamei K, Goldman GH. 2016. Epidemiological and genomic landscape of azole resistance mechanisms in *Aspergillus* fungi. Front Microbiol 7:1382. <https://doi.org/10.3389/fmicb.2016.01382>.
  145. Hodiament CJ, Dolman KM, Ten Berge IJ, Melchers WJ, Verweij PE, Pajkrt D. 2009. Multiple-azole-resistant *Aspergillus fumigatus* osteomyelitis in a patient with chronic granulomatous disease successfully

- treated with long-term oral posaconazole and surgery. *Med Mycol* 47:217–220. <https://doi.org/10.1080/13693780802545600>.
146. Paul S, Klutts JS, Moye-Rowley WS. 2012. Analysis of promoter function in *Aspergillus fumigatus*. *Eukaryot Cell* 11:1167–1177. <https://doi.org/10.1128/EC.00174-12>.
  147. Kato M. 2005. An overview of the CCAAT-box binding factor in filamentous fungi: assembly, nuclear translocation, and transcriptional enhancement. *Biosci Biotechnol Biochem* 69:663–672. <https://doi.org/10.1271/bbb.69.663>.
  148. Brakhage AA, Andrianopoulos A, Kato M, Steidl S, Davis MA, Tsukagoshi N, Hynes MJ. 1999. HAP-like CCAAT-binding complexes in filamentous fungi: implications for biotechnology. *Fungal Genet Biol* 27:243–252. <https://doi.org/10.1006/fgbi.1999.1136>.
  149. Paul RA, Rudramurthy SM, Meis JF, Mouton JW, Chakrabarti A. 2015. A novel Y319H substitution in CYP51C associated with azole resistance in *Aspergillus flavus*. *Antimicrob Agents Chemother* 59:6615–6619. <https://doi.org/10.1128/AAC.00637-15>.
  150. Zhao Y, Garnaud C, Brenier-Pinchart MP, Thiebaut-Bertrand A, Saint-Raymond C, Camara B, Hamidfar R, Cognet O, Maubon D, Cornet M, Perlin DS. 2016. Direct molecular diagnosis of aspergillosis and CYP51A profiling from respiratory samples of French patients. *Front Microbiol* 7:1164. <https://doi.org/10.3389/fmicb.2016.01164>.
  151. Araujo R, Gungor O, Amorim A. 2015. Single-tube PCR coupled with mini-sequencing assay for the detection of *cyp51A* and *cyp51B* polymorphisms in *Aspergillus fumigatus*. *Future Microbiol* 10:1797–1804. <https://doi.org/10.2217/fmb.15.103>.
  152. Liu M, Zheng N, Li D, Zheng H, Zhang L, Ge H, Liu W. 2016. *cyp51A*-based mechanism of azole resistance in *Aspergillus fumigatus*: illustration by a new 3D structural model of *Aspergillus fumigatus* CYP51A protein. *Med Mycol* 54:400–408. <https://doi.org/10.1093/mmy/myv102>.
  153. O’Gorman CM, Fuller H, Dyer PS. 2009. Discovery of a sexual cycle in the opportunistic fungal pathogen *Aspergillus fumigatus*. *Nature* 457:471–474. <https://doi.org/10.1038/nature07528>.
  154. Abdolrasouli A, Rhodes J, Beale MA, Hagen F, Rogers TR, Chowdhary A, Meis JF, Armstrong-James D, Fisher MC. 2015. Genomic context of azole resistance mutations in *Aspergillus fumigatus* determined using whole-genome sequencing. *mBio* 6:e00536-14. <https://doi.org/10.1128/mBio.00536-15>.
  155. Tashiro M, Izumikawa K, Hirano K, Ide S, Mihara T, Hosogaya N, Takazono T, Morinaga Y, Nakamura S, Kurihara S, Imamura Y, Miyazaki T, Nishino T, Tsukamoto M, Takeya H, Yamamoto Y, Yanagihara K, Yasuoka A, Tashiro T, Kohno S. 2012. Correlation between triazole treatment history and susceptibility in clinically isolated *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 56:4870–4875. <https://doi.org/10.1128/AAC.00514-12>.
  156. Alcazar-Fuoli L, Mellado E, Cuenca-Estrella M, Sanglard D. 2011. Probing the role of point mutations in the *cyp51A* gene from *Aspergillus fumigatus* in the model yeast *Saccharomyces cerevisiae*. *Med Mycol* 49:276–284. <https://doi.org/10.3109/13693786.2010.512926>.
  157. Lescar J, Meyer I, Akshita K, Srinivasaraghavan K, Verma C, Palous M, Mazier D, Datry A, Fekkar A. 2014. *Aspergillus fumigatus* harbouring the sole Y121F mutation shows decreased susceptibility to voriconazole but maintained susceptibility to itraconazole and posaconazole. *J Antimicrob Chemother* 69:3244–3247. <https://doi.org/10.1093/jac/dku316>.
  158. Snelders E, Camps SM, Karawajczyk A, Rijs AJ, Zoll J, Verweij PE, Melchers WJ. 2015. Genotype-phenotype complexity of the TR<sub>46</sub>/Y121F/T289A *cyp51A* azole resistance mechanism in *Aspergillus fumigatus*. *Fungal Genet Biol* 82:129–135. <https://doi.org/10.1016/j.fgb.2015.06.001>.
  159. Belleste B, Raberin H, Morel J, Flori P, Hafid J, Manhungsung RT. 2010. Acquired resistance to voriconazole and itraconazole in a patient with pulmonary aspergilloma. *Med Mycol* 48:197–200. <https://doi.org/10.3109/13693780902717018>.
  160. Krishnan-Natesan S, Chandrasekar PH, Alangaden GJ, Manavathu EK. 2008. Molecular characterisation of *cyp51A* and *cyp51B* genes coding for P450 14 $\alpha$ -lanosterol demethylases A (CYP51Ap) and B (CYP51Bp) from voriconazole-resistant laboratory isolates of *Aspergillus flavus*. *Int J Antimicrob Agents* 32:519–524. <https://doi.org/10.1016/j.ijantimicag.2008.06.018>.
  161. Sharma C, Kumar R, Kathuria S, Kumar Singh P, Gupta D, Chowdhary A. 2016. Prevalence of azole resistance in clinical *Aspergillus flavus* isolates: presence of novel S196F, A324P, N423D and V465M substitutions in the *cyp51C* gene, abstr 4352. 26th Eur Congr Clin Microbiol Infect Dis Conf Abstr.
  162. Liu W, Sun Y, Chen W, Liu W, Wan Z, Bu D, Li R. 2012. The T788G mutation in the *cyp51C* gene confers voriconazole resistance in *Aspergillus flavus* causing aspergillosis. *Antimicrob Agents Chemother* 56:2598–2603. <https://doi.org/10.1128/AAC.05477-11>.
  163. Nierman WC, Yu J, Fedorova-Abrams ND, Losada L, Cleveland TE, Bhatnagar D, Bennett JW, Dean R, Payne GA. 2015. Genome sequence of *Aspergillus flavus* NRRL 3357, a strain that causes aflatoxin contamination of food and feed. *Genome Announc* 3:e00168-15. <https://doi.org/10.1128/genomeA.00168-15>.
  164. Rudramurthy SM, Seyedmousavi S, Dhaliwal M, Chakrabarti A, Meis JF, Mouton JW. 7 November 2016. Pharmacodynamics of voriconazole against wild-type and azole-resistant *Aspergillus flavus* isolates in a non-neutropenic murine model of disseminated aspergillosis. *Antimicrob Agents Chemother* <https://doi.org/10.1128/AAC.01491-16>.
  165. Arendrup MC, Jensen RH, Grif K, Skov M, Pressler T, Johansen HK, Lass-Flörl C. 2012. *In vivo* emergence of *Aspergillus terreus* with reduced azole susceptibility and a Cyp51A M217I alteration. *J Infect Dis* 206:981–985. <https://doi.org/10.1093/infdis/jis442>.
  166. Iatta R, Nuccio F, Immediato D, Mosca A, De Carlo C, Miragliotta G, Parisi A, Crescenzo G, Otranto D, Cafarchia C. 2016. Species distribution and *in vitro* azole susceptibility of *Aspergillus* section Nigri isolates from clinical and environmental settings. *J Clin Microbiol* 54:2365–2372. <https://doi.org/10.1128/JCM.01075-16>.
  167. Ahmad S, Khan Z, Hagen F, Meis JF. 2014. Simple, low-cost molecular assays for TR<sub>34</sub>/L98H mutations in the *cyp51A* gene for rapid detection of triazole-resistant *Aspergillus fumigatus* isolates. *J Clin Microbiol* 52:2223–2227. <https://doi.org/10.1128/JCM.00408-14>.
  168. Howard SJ, Arendrup MC. 2011. Acquired antifungal drug resistance in *Aspergillus fumigatus*: epidemiology and detection. *Med Mycol* 49(Suppl 1):S90–S95. <https://doi.org/10.3109/13693786.2010.508469>.
  169. Chen J, Li H, Li R, Bu D, Wan Z. 2005. Mutations in the *cyp51A* gene and susceptibility to itraconazole in *Aspergillus fumigatus* serially isolated from a patient with lung aspergilloma. *J Antimicrob Chemother* 55:31–37. <https://doi.org/10.1093/jac/dkh507>.
  170. Camps SM, Rijs AJ, Klaassen CH, Meis JF, O’Gorman CM, Dyer PS, Melchers WJ, Verweij PE. 2012. Molecular epidemiology of *Aspergillus fumigatus* isolates harboring the TR<sub>34</sub>/L98H azole resistance mechanism. *J Clin Microbiol* 50:2674–2680. <https://doi.org/10.1128/JCM.00335-12>.
  171. Montesinos I, Dodemont M, Lagrou K, Jacobs F, Etienne I, Denis O. 2014. New case of azole-resistant *Aspergillus fumigatus* due to TR<sub>46</sub>/Y121F/T289A mutation in Belgium. *J Antimicrob Chemother* 69:3439–3440. <https://doi.org/10.1093/jac/dku289>.
  172. Weber M, Schaer J, Walther G, Kaerger K, Steinmann J, Rath PM, Spiess B, Buchheidt D, Hamprecht A, Kurzai O. 11 March 2017. FunResDB—a web resource for genotypic susceptibility testing of *Aspergillus fumigatus*. *Med Mycol* <https://doi.org/10.1093/mmy/myx015>.
  173. Koehler P, Hamprecht A, Bader O, Bekeredjian-Ding I, Buchheidt D, Doelken G, Elias J, Haase G, Hahn-Ast C, Karthaus M, Kekule A, Keller P, Kiehl M, Krause SW, Kramer C, Neumann S, Rohde H, La Rosee P, Ruhnke M, Schafhausen P, Schalk E, Schulz K, Schwartz S, Silling G, Staib P, Ullmann A, Vergoulidou M, Weber T, Cornely OA, Vehreschild MJ. 2017. Epidemiology of invasive aspergillosis and azole resistance in patients with acute leukaemia: the SEPIA Study. *Int J Antimicrob Agents* 49:218–223. <https://doi.org/10.1016/j.ijantimicag.2016.10.019>.
  174. Spiess B, Postina P, Reinwald M, Cornely OA, Hamprecht A, Hoenigl M, Lass-Flörl C, Rath PM, Steinmann J, Miethke T, Lauten M, Will S, Merker N, Hofmann WK, Buchheidt D. 2014. Incidence of *cyp51A* key mutations in *Aspergillus fumigatus*—a study on primary clinical samples of immunocompromised patients in the period of 1995–2013. *PLoS One* 9:e103113. <https://doi.org/10.1371/journal.pone.0103113>.
  175. Spiess B, Seifarth W, Merker N, Howard SJ, Reinwald M, Dietz A, Hofmann WK, Buchheidt D. 2012. Development of novel PCR assays to detect azole resistance-mediating mutations of the *Aspergillus fumigatus* *cyp51A* gene in primary clinical samples from neutropenic patients. *Antimicrob Agents Chemother* 56:3905–3910. <https://doi.org/10.1128/AAC.05902-11>.
  176. Chong GL, van de Sande WW, Dingemans GJ, Gaajetaan GR, Vonk AG, Hayette MP, van Tegelen DW, Simons GF, Rijnders BJ. 2015. Validation of a new *Aspergillus* real-time PCR assay for direct detection of *Aspergillus* and azole resistance of *Aspergillus fumigatus* on bronchoalveolar

- lavage fluid. *J Clin Microbiol* 53:868–874. <https://doi.org/10.1128/JCM.03216-14>.
177. White PL, Posso RB, Barnes RA. 2015. Analytical and clinical evaluation of the PathoNostics AsperGenius assay for detection of invasive aspergillosis and resistance to azole antifungal drugs during testing of serum samples. *J Clin Microbiol* 53:2115–2121. <https://doi.org/10.1128/JCM.00667-15>.
  178. Coste A, Turner V, Ischer F, Morschhauser J, Forche A, Selmecki A, Berman J, Bille J, Sanglard D. 2006. A mutation in Tac1p, a transcription factor regulating *CDR1* and *CDR2*, is coupled with loss of heterozygosity at chromosome 5 to mediate antifungal resistance in *Candida albicans*. *Genetics* 172:2139–2156. <https://doi.org/10.1534/genetics.105.054767>.
  179. Hagiwara D, Miura D, Shimizu K, Paul S, Ohba A, Gono T, Watanabe A, Kamei K, Shintani T, Moye-Rowley WS, Kawamoto S, Gomi K. 2017. A novel Zn2-Cys6 transcription factor AtrR plays a key role in an azole resistance mechanism of *Aspergillus fumigatus* by co-regulating *cyp51A* and *cdr1B* expressions. *PLoS Pathog* 13:e1006096. <https://doi.org/10.1371/journal.ppat.1006096>.
  180. Blatzer M, Barker BM, Willger SD, Beckmann N, Blosser SJ, Cornish EJ, Mazurie A, Grahl N, Haas H, Cramer RA. 2011. SREBP coordinates iron and ergosterol homeostasis to mediate triazole drug and hypoxia responses in the human fungal pathogen *Aspergillus fumigatus*. *PLoS Genet* 7:e1002374. <https://doi.org/10.1371/journal.pgen.1002374>.
  181. Lamoth F, Juvvadi PR, Soderblom EJ, Moseley MA, Asfaw YG, Steinbach WJ. 2014. Identification of a key lysine residue in heat shock protein 90 required for azole and echinocandin resistance in *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 58:1889–1896. <https://doi.org/10.1128/AAC.02286-13>.
  182. Losada L, Sugui JA, Eckhaus MA, Chang YC, Mounaud S, Figat A, Joardar V, Pakala SB, Pakala S, Venepally P, Fedorova N, Nierman WC, Kwon-Chung KJ. 2015. Genetic analysis using an isogenic mating pair of *Aspergillus fumigatus* identifies azole resistance genes and lack of MAT locus's role in virulence. *PLoS Pathog* 11:e1004834. <https://doi.org/10.1371/journal.ppat.1004834>.
  183. de Valk HA, Meis JF, Curfs IM, Muehlethaler K, Mouton JW, Klaassen CH. 2005. Use of a novel panel of nine short tandem repeats for exact and high-resolution fingerprinting of *Aspergillus fumigatus* isolates. *J Clin Microbiol* 43:4112–4120. <https://doi.org/10.1128/JCM.43.8.4112-4120.2005>.
  184. Gao LJ, Sun Y, Wan Z, Li RY, Yu J. 2013. CSP typing of Chinese *Aspergillus fumigatus* isolates: identification of additional CSP types. *Med Mycol* 51:683–687. <https://doi.org/10.3109/13693786.2013.770609>.
  185. Kidd SE, Nik Zulkepli NA, Slavin MA, Morrissey CO. 2009. Utility of a proposed CSP typing nomenclature for Australian *Aspergillus fumigatus* isolates: identification of additional CSP types and suggested modifications. *J Microbiol Methods* 78:223–226. <https://doi.org/10.1016/j.mimet.2009.06.003>.
  186. Klaassen CH, de Valk HA, Balajee SA, Meis JF. 2009. Utility of CSP typing to sub-type clinical *Aspergillus fumigatus* isolates and proposal for a new CSP type nomenclature. *J Microbiol Methods* 77:292–296. <https://doi.org/10.1016/j.mimet.2009.03.004>.
  187. Garcia-Rubio R, Gil H, Monteiro MC, Pelaez T, Mellado E. 2016. A new *Aspergillus fumigatus* typing method based on hypervariable tandem repeats located within exons of surface protein coding genes (TRESP). *PLoS One* 11:e0163869. <https://doi.org/10.1371/journal.pone.0163869>.
  188. Ashu EE, Hagen F, Chowdhary A, Meis JF, Xu J. 2017. Global population genetic analysis of *Aspergillus fumigatus*. *mSphere* 2:e00019-17. <https://doi.org/10.1128/mSphere.00019-17>.
  189. Valsecchi I, Mellado E, Beau R, Raj S, Latge JP. 2015. Fitness studies of azole-resistant strains of *Aspergillus fumigatus*. *Antimicrob Agents Chemother* 59:7866–7869. <https://doi.org/10.1128/AAC.01594-15>.
  190. Vazquez JA, Manavathu EK. 2016. Molecular characterization of a voriconazole-resistant, posaconazole-susceptible *Aspergillus fumigatus* isolate in a lung transplant recipient in the United States. *Antimicrob Agents Chemother* 60:1129–1133. <https://doi.org/10.1128/AAC.01130-15>.
  191. Astvad KMT, Hare RK, Arendrup MC. 2017. Evaluation of the in vitro activity of isavuconazole and comparator voriconazole against 2635 contemporary clinical *Candida* and *Aspergillus* isolates. *Clin Microbiol Infect* 2017:S1198-743X(17)30192-1. <https://doi.org/10.1016/j.cmi.2017.03.023>.
  192. van der Linden JW, Snelders E, Arends JP, Daenen SM, Melchers WJ, Verweij PE. 2010. Rapid diagnosis of azole-resistant aspergillosis by direct PCR using tissue specimens. *J Clin Microbiol* 48:1478–1480. <https://doi.org/10.1128/JCM.02221-09>.
  193. Denning DW, Park S, Lass-Flörl C, Fraczek MG, Kirwan M, Gore R, Smith J, Bueid A, Moore CB, Bowyer P, Perlin DS. 2011. High-frequency triazole resistance found in nonculturable *Aspergillus fumigatus* from lungs of patients with chronic fungal disease. *Clin Infect Dis* 52:1123–1129. <https://doi.org/10.1093/cid/cir179>.
  194. Klaassen CH, de Valk HA, Curfs-Breuker IM, Meis JF. 2010. Novel mixed-format real-time PCR assay to detect mutations conferring resistance to triazoles in *Aspergillus fumigatus* and prevalence of multi-triazole resistance among clinical isolates in the Netherlands. *J Antimicrob Chemother* 65:901–905. <https://doi.org/10.1093/jac/dkq041>.

**Anna Dudakova** is a medical microbiologist with a focus on antimicrobial stewardship, including antifungal treatment. Infections with *Aspergillus* spp. and *Candida* spp. are her main interest. She works at the Institute for Medical Microbiology at the University Medical Center Göttingen, Germany.



**Birgit Spiess** is a molecular biologist with a research focus on molecular diagnostics of invasive fungal diseases and azole resistance in *Aspergillus fumigatus*, especially in patients with hematological malignancies. She works as a Senior Scientist in the Scientific Laboratory of the Department of Hematology and Oncology at the Mannheim University Hospital, Heidelberg University, Germany.



**Marut Tangwattanachuleeporn** is a microbiologist currently holding an Assistant Professorship for Microbiology at Burapha University, Chon Buri, Thailand. His studies focus on the prevalence of pathogenic fungi in the environment, such as *Aspergillus* spp., dermatophytes, and *Cryptococcus* spp., as potential sources of infection, as well as monitoring of their antimycotic drug resistance.




**Christoph Sasse** is a molecular microbiologist with a research focus on the stress and drug responses of the pathogenic fungus *Aspergillus fumigatus*. He was trained in Göttingen, Germany, at Georg August University and currently holds a postdoctoral position in the Department of Molecular Microbiology and Genetics of Georg August University.





**Dieter Buchheidt** is a hematologist and infectious diseases specialist with a focus on biomarker-based diagnosis of *Aspergillus* infections in patients with hematological malignancies. He is a Senior Physician at the Mannheim University Hospital, University of Heidelberg, Germany.



**Michael Weig** is a medical microbiologist with a research focus on the pathogenesis, epidemiology, diagnosis, and therapy of human fungal infections. He is a Senior Physician at the Institute for Medical Microbiology at the University Medical Center Göttingen.



**Uwe Groß** is a medical microbiologist who is focusing his research activities on the epidemiology and diagnosis of infectious diseases. He is Head of the Institute for Medical Microbiology at the University Medical Center Göttingen.



**Oliver Bader** is a molecular biologist with a research focus on pathogenic fungi. Initially trained in molecular mycology at the Robert Koch Institute, he is currently a group leader at the Institute for Medical Microbiology, working on the development and refinement of diagnostic strategies, including susceptibility testing, MALDI-TOF mass spectrometry, and serologic detection of bacterial and fungal infections.

